

ZETA CORRESPONDENCES IN RANK- n

A. ÁLVAREZ*

CONTENTS

1. Introduction.	1
2. Vector bundles with level structures and zeta correspondences over $\mathbb{P}_1$	4
2.1. Vector bundles with level structures	4
2.2. Zeta correspondences for vector bundles over $\mathbb{P}_1$ with level structures	6
3. Drinfeld modules and zeta correspondences	10
3.1. A -Drinfeld modules, elliptic sheaves and their antiequivalence	10
3.2. A -Drinfeld modules and elliptic sheaves with level structures	13
3.3. $\mathbb{F}_q[t]$ -Drinfeld modules and zeta correspondences	15
3.4. Some explicit irreducible components for zeta correspondences	18
3.5. Examples	20
4. Case of a general curve	24
4.1. Zeta subschemes for rank- n -Shtukas	29
4.2. Zeta subschemes and A -Drinfeld modules	31
References	35

1. INTRODUCTION.

The purpose of this work is to study in the case of function fields and rank- n certain correspondences associated with Jacobi sums [C]. These correspondences are called zeta correspondences and have been widely studied [An2], [An3], [An4], [AnDP], [Si]. Their importance is the relationship with Stickelberger's Theorem and L -series evaluators, [An2], [An3], [An4], [AnDP]. In work [C] they are used to prove that the Frobenius map is essentially a Gauss or Jacobi sum for Fermat and Artin-Schreier curves.

In rank 1, these zeta correspondences are closely related to the theta divisor defined over the Jacobian, [An2]. The strategy in works [An2], [An3] is to study this subject in the setting of Drinfeld modules. Here we shall follow the same strategy but considering $\mathbb{F}_q[t]$ -Drinfeld modules (elliptic sheaves) with level structures over an effective divisor D on $\text{Spec}(\mathbb{F}_q[t])$. In this way, the moduli of Drinfeld modules (elliptic sheaves) with level structures are given by the coefficients of a polynomial. These coefficients are torsion elements for a universal Drinfeld module (c.f [Al]): In this setting, zeta correspondences are given explicitly in terms of these elements. To study rank- n zeta correspondences, we have taken into account the generalized theta function considered in [An3] and the morphism given in 4. [An2], but considering

MSC: 11G09, 14D20, 14K25

* Departamento de Matemáticas. Universidad de Salamanca. Spain.

the generalized genus [Se] pg.86. Firstly, we shall introduce zeta subschemes over the moduli of vector bundles with level structures as a generalization of the n^2 -generalized theta divisor. We recall that the n^2 -generalized theta divisor over a curve C of genus g is formed by the vector bundles of rank n^2 and degree $n^2(g-1)$ without global sections. The restriction via a certain morphism, similar to that of 4. [An2], of these zeta subschemes to the moduli of rank- n Drinfeld modules (elliptic sheaves) will be the zeta correspondences that we wish to study.

In this way we shall explicitly obtain, for $\mathbb{P}_1$, the zeta correspondences as the zero locus of certain functions. In the rank- n case, they are the zero locus of n^2 functions given by a $n \times n$ -matrix. These functions involve the torsion elements of a rank- n universal $\mathbb{F}_q[t]$ -Drinfeld module. To obtain these results we shall prove a similar Lemma to " $\chi = 0 \Rightarrow h^0 = h^1 = 0$ " Lemma 3.3.1, [An2] for level structures in the rank- n case and $\mathbb{P}_1$. We can prove this Lemma in our case because for semistable bundles over $\mathbb{P}_1$ the generalized theta divisor is the empty subset. For rank-1, as in [An2], we shall see a relationship of these zeta correspondences with the theta divisor for level structures. One observes that instead of the zero locus of the $n \times n$ -matrix considered one could take the zero locus of its determinant and would thus obtain a 1-correspondence that involves the zeta correspondence considered before. The examples that we provide in the rank-2 case give us zeta correspondences whose irreducible components are essentially the same as in the rank-1 case. They are given by graphs of automorphisms given by central matrices and the Frobenius morphism.

Above we have considered Drinfeld modules with level structures over $\infty + D$, ∞ being the pole of the Drinfeld modules. However if we consider another rational point p different from ∞ , we can extend the above results to an arbitrary proper and smooth curve C over a finite field. We shall obtain zeta subschemes for rank- n shtukas; these zeta subschemes would be formed by pairs of isogenous shtukas although to complete the result it would be necessary to prove the " $\chi = 0 \Rightarrow h^0 = h^1 = 0$ "-Lemma for rank n . This result tells us, via the immersion of A -Drinfeld modules in rank- n shtukas, that for $\mathbb{P}_1$ zeta correspondences (p instead of ∞) are given by pairs of isogenous A -Drinfeld modules.

In the case of a general curve, these zeta subschemes are again given, now locally, by the zero locus of n^2 -functions. We shall check that these functions are the restriction of hyperplane sections via a Plucker morphism. In this way, these zeta subschemes are the intersection of n^2 -generalized theta divisors.

I believe that it would be very interesting to complete section 4.1 in the setting of the iterated shtukas and Lafforgue's compactification for the shtuka variety [La1], [La2] and the arithmetic counterpart.

The paper is organized in the following way:

Section 2 is devoted to recalling some definitions and results of rank- n vector bundles with level structures. In subsection 2.2, we introduce zeta correspondences for the moduli of semistable vector bundles with level structures over $\mathbb{P}_1$, which are analogous to the n^2 -generalized theta divisor. Moreover, zeta correspondences are given by the zero locus of n^2 -functions, obtained from a matrix.

In section 3. we recall some definitions and results for rank- n Drinfeld modules and elliptic sheaves with level structures. Via the immersion of elliptic sheaves in the moduli of semistable vector bundles, we gain a notion of zeta correspondences for rank- n Drinfeld modules. By means of a similar Lemma to Lemma 3.3.1 [An2],

we can give some information about the irreducible components of these zeta correspondences. We provide rank-2 examples and the known rank-1 examples, [An3], [AnDP], [Si].

In section 4, by taking a rational point p instead of ∞ , we can obtain in a more precise way the results of the last sections for a general curve C and rank- n shtukas. We state a similar Lemma to Lemma 3.3.1 [An2] for rank- n shtukas, although in this case we should establish this Lemma in rank n without level structures to complete this Lemma.

List of notations and previous definitions

$\mathbb{F}_q$ is a finite field with q -elements, ($q = p^m$)

$\otimes$ denotes $\otimes_{\mathbb{F}_q}$

R is a $\mathbb{F}_q$ -algebra

C is a smooth, proper and geometrically irreducible curve over $\mathbb{F}_q$

g is the genus of C

∞ and p are rational points in C

$A = H^0(C \setminus \{\infty\}, \mathcal{O}_C)$

D is an effective rational divisor over $\text{Spec}(A)$

If M is a vector bundle over C , $M(k)$ denotes $M \otimes_{\mathcal{O}_C} \mathcal{O}_C(k\infty)$, $k \in \mathbb{Z}$.

$R^\times$ denotes the group of units in a ring R

If $s \in \text{Spec}(R)$, we denote by m_s the prime ideal associated with s .

If M is a vector bundle over $C \times \text{Spec}(R)$, and $s \in \text{Spec}(R)$, $\deg(M_s)$ denotes the degree of the vector bundle, M_s , over $C \times k(s)$. In this paper we shall consider vector bundles of constant degree, $\deg(M)$, for each $s \in \text{Spec}(R)$.

Let M be a vector bundle over $\mathcal{O}_C$; $M^\vee$ denotes the dual vector bundle

$(\mathbb{G}_a)_R$ is the additive line group over a ring R

F denotes the Frobenius morphism over a scheme $\text{Spec}(R)$

For a vector bundle, M , over $C \times \text{Spec}(R)$, $F^\#(M)$ denotes the pull-back $(Id \times F)^*M$.

If $T : M \rightarrow M'$ is a morphism of vector bundles over a scheme, $H^0(T)$ denotes the morphism induced among the global sections

2. VECTOR BUNDLES WITH LEVEL STRUCTURES AND ZETA CORRESPONDENCES OVER $\mathbb{P}_1$

2.1. Vector bundles with level structures.

Definition 2.1. A ∞D -level structure is a pair $(M, f_{\infty D})$, where M is a rank- n vector bundle over $C \times \text{Spec}(R)$ and

$$f_{\infty D} : M \rightarrow (\mathcal{O}_C/\mathcal{O}_C(-\infty - D) \otimes R)^n$$

is a surjective morphism of $\mathcal{O}_C \otimes R$ -modules.

By $\mathcal{O}_C/\mathcal{O}_C(-\infty - D)$ and $\mathcal{O}_{\infty D}$ we denote $(\mathcal{O}_{\infty D})$ and $H^0(C, (\mathcal{O}_C/\mathcal{O}_C(-\infty - D)))$ respectively.

A morphism of two pairs, $(M, f_{\infty D})$, $(M', f'_{\infty D})$, is a morphism of $\mathcal{O}_C \otimes R$ -modules, $\phi : M \rightarrow M'$, such that the diagram

$$\begin{array}{ccc} M & \xrightarrow{\phi} & M' \\ & \searrow \bar{f}_{\infty D} & \downarrow f'_{\infty D} \\ & & (\mathcal{O}_{-\infty - D})^n \otimes R \end{array}$$

is commutative. Two pairs $(M, f_{\infty D})$, $(M', f'_{\infty D})$ are said to be equivalent if there exists an isomorphism ϕ .

Proposition 2.2. If $\phi : M \rightarrow M'$ is a morphism between two pairs $(M, f_{\infty D})$, $(M', f'_{\infty D})$ defined over a field K , then ϕ is injective.

Proof. Since $M/\text{Ker}(\phi) \hookrightarrow M'$, $M/\text{Ker}(\phi)$ is a free torsion module, and since

$$M/\text{Ker}(\phi) \otimes_{\mathcal{O}_C} \mathcal{O}_{\infty D} \simeq (\mathcal{O}_{\infty D})^n \otimes K,$$

$M/\text{Ker}(\phi)$ is locally free of rank n , and hence $\text{Ker}(\phi) = 0$ □

We recall that M is said to be semistable if for each geometric point $s \in \text{Spec}(R)$, $M_s := M \otimes_R k(s)$ is semistable; i.e, for each $F \subset M_s$

$$\mu(F) := \deg(F)/\text{rank}(F) \leq \mu(M_s)$$

Let denote us by $\mathcal{M}_C^{ss}(n, h, \infty D)$ the moduli scheme of pairs $(M, f_{\infty D})$ of semistable vector bundles, M , of rank n , degree h , with a ∞D -level structure, over a curve C (c.f.:[Ss]). Then,

$$\text{Hom}_{\text{Schem.}}(\text{Spec}(R), \mathcal{M}_C^{ss}(n, h, \infty D)) = \{ \text{pairs over } R, (M, f_{\infty D}) \} / \text{Up to isomor.}$$

In the same way, we shall denote $\mathcal{M}_C(n, h, \infty D)$, the moduli stack, [LM], of pairs $(M, f_{\infty D})$, of vector bundles, M , of rank n , degree h , with a ∞D -level structure.

In all this section, $C = \mathbb{P}_1$. By choosing a local parameter $1/t$ in ∞ , $A = \mathbb{F}_q[t]$. We consider an isomorphism $H^0(\mathcal{O}_{\mathbb{P}_1}, \mathcal{O}_{\mathbb{P}_1}/\mathcal{O}_{\mathbb{P}_1}(-\infty - D)) = \mathbb{F}_q[t]/p(t) \times \mathbb{F}_q$, with $p(t) = (t - \alpha_1)^{r_1} \cdot (t - \alpha_2)^{r_2} \cdots (t - \alpha_l)^{r_l}$ and $D = r_1.x_1 + \cdots + r_l.x_l$, where x_i is the point associated with the maximal $(t - \alpha_i)\mathbb{F}_q[t]$. Recall that $d = \deg(D)$.

Proposition 2.3. Let M be a semistable vector bundle over $\mathbb{P}_1 \times \text{Spec}(R)$ of rank n and degree 0 with a ∞D -level structure $f_{\infty D}$. Then:

1) $H^0(\mathbb{P}_1 \times \text{Spec}(R), M)$ is a free R -module of rank n , and $M = H^0(\mathbb{P}_1 \times \text{Spec}(R), M) \otimes \mathcal{O}_C$.

2) Two pairs $(M, f_{\infty D})$, $(\bar{M}, \bar{f}_{\infty D})$, $\deg(M) = \deg(\bar{M}) = 0$, are equivalent if and only if by choosing bases $\{s_1, \dots, s_n\}$ and $\{\bar{s}_1, \dots, \bar{s}_n\}$ for $H^0(\mathbb{P}_1 \times \text{Spec}(R), M)$ and $H^0(\mathbb{P}_1 \times \text{Spec}(R), \bar{M})$ respectively, there exists $g \in \text{Gl}_n(R)$, with

$$f_{\infty D}(s_i) = \bar{f}_{\infty D}(g(\bar{s}_i))$$

Proof. 1) By taking global sections in the exact sequence of $\mathcal{O}_{\mathbb{P}_1} \otimes R$ -modules

$$0 \rightarrow M(-\infty) \rightarrow M \rightarrow M/M(-\infty) \rightarrow 0,$$

and bearing in mind the morphism given by the D -level structure $f_{\infty} : M \rightarrow (\mathcal{O}_{\infty} \otimes R)^n$, we obtain an isomorphism

$$M/M(-\infty) \simeq (\mathcal{O}_{\infty} \otimes R)^n$$

and we conclude because, as M is semistable, we have

$$H^0(\mathbb{P}_1 \times \text{Spec}(R), M(-\infty)) = H^1(\mathbb{P}_1 \times \text{Spec}(R), M(-\infty)) = 0$$

and therefore $H^0(f_{\infty}) : H^0(\mathbb{P}_1 \times \text{Spec}(R), M) \rightarrow (\mathcal{O}_{\infty} \otimes R)^n$ is an isomorphism. The second assertion is easily deduced because $H^0(\mathbb{P}_1 \times \text{Spec}(R), M) \otimes \mathcal{O}_C \subseteq M$ and $\deg(M) = 0$.

2) If $(M, f_{\infty D})$, $(\bar{M}, \bar{f}_{\infty D})$ are equivalent, there exists an isomorphism of modules $\phi : M \rightarrow \bar{M}$ such that $\bar{f}_{\infty D} \cdot \phi = f_{\infty D}$. In this case, $g \in \text{Gl}_n(R)$ is given by considering $H^0(\phi)$ and bases for $H^0(\mathbb{P}_1 \times \text{Spec}(R), M)$ and $H^0(\mathbb{P}_1 \times \text{Spec}(R), \bar{M})$.

Conversely, if $\{s_1, \dots, s_n\}$ and $\{\bar{s}_1, \dots, \bar{s}_n\}$ are bases for $H^0(\mathbb{P}_1 \times \text{Spec}(R), M)$ and $H^0(\mathbb{P}_1 \times \text{Spec}(R), \bar{M})$ respectively, satisfying

$$f_{\infty D}(s_i) = \bar{f}_{\infty D}(g(\bar{s}_i))$$

for all i , ϕ is obtained from g , since g can be interpreted as an isomorphism:

$$M \simeq (\mathcal{O}_{\mathbb{P}_1} \otimes R).s_1, \dots, (\mathcal{O}_{\mathbb{P}_1} \otimes R).s_n \rightarrow \bar{M} \simeq (\mathcal{O}_{\mathbb{P}_1} \otimes R).\bar{s}_1, \dots, (\mathcal{O}_{\mathbb{P}_1} \otimes R).\bar{s}_n$$

□

Remark 1. By considering a basis $\{s_1, \dots, s_n\}$, by the last Lemma $(M, f_{\infty D})$ has associated, in a bijective way, a matrix:

$$(\Delta_0 + \Delta_1 t + \dots + \Delta_{d-1} t^{d-1}) \times \Delta_{\infty}$$

where $\Delta_k, \Delta_{\infty}$ are $n \times n$ -matrices with entry elements in R . These matrices are given by $\{f_{\infty D}(s_1), \dots, f_{\infty D}(s_n)\} \subset (\mathcal{O}_{\infty D})^n \otimes R$.

One can consider the vector bundle (scheme) over $\mathbb{F}_q$, $V(\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (\mathcal{O}_{\infty D})^n))$ associated with the $\mathbb{F}_q$ -vector space $\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (\mathcal{O}_{\infty D})^n)$. Then, for this vector bundle there exists a universal object

$$(\mathcal{D}_0 + \mathcal{D}_1 t + \dots + \mathcal{D}_{d-1} t^{d-1}) \times \mathcal{D}_{\infty}$$

$\mathcal{D}_k$ and $\mathcal{D}_{\infty}$ being matrices with the independent variables $d_{i,j,k}$ and $d_{i,j,\infty}$ as entries, respectively.

Lemma 2.4. There exists an injective immersion (of schemes)

$$\Psi : \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D) \longrightarrow V(\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (O_{\infty D})^n))/Gl_n$$

where Gl_n denotes the n -linear algebraic group $/\mathbb{F}_q$ acting on

$$V(\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (O_{\infty D})^n))$$

over the term $\mathbb{F}_q^n$.

Proof. Noticing that $V(\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (O_{\infty D})^n))/Gl_n$ is the Grassmannian of n -planes in $(O_{\infty D})^n$, then $\Psi(M, f_{\infty D})$ is defined as the rank- n subbundle of $(O_{\infty D})^n$ generated by $\{f_{\infty D}(s_1), \dots, f_{\infty D}(s_n)\}$. If

$$(M, f_{\infty D}) \in \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D)(R),$$

then $\langle f_{\infty D}(s_1), \dots, f_{\infty D}(s_n) \rangle$ is a rank- n subbundle of $(O_{\infty D})^n \otimes R$, because taking sections in the exact sequence

$$0 \rightarrow M(-\infty - D) \rightarrow M \xrightarrow{f_{\infty D}} (O_{\infty D})^n \otimes R \rightarrow 0$$

one obtains the exact sequence

$$\langle f_{\infty D}(s_1), \dots, f_{\infty D}(s_n) \rangle \hookrightarrow (O_{\infty D})^n \otimes R \rightarrow H^1(\mathbb{P}_1 \times \text{Spec}(R), M(-\infty - D)) \rightarrow 0.$$

To obtain this last exact sequence we have used the fact that M is semistable and hence $H^1(\mathbb{P}_1 \times \text{Spec}(R), M) = 0$ and $H^0(\mathbb{P}_1 \times \text{Spec}(R), M(-\infty - D)) = 0$. Moreover, by Grauert's theorem $H^1(\mathbb{P}_1 \times \text{Spec}(R), M(-\infty - D))$ is a locally free R -module. $\square$

Remark 2. Ψ takes values in the open subscheme, U , of the vector bundle of

$$\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (O_D)^n)$$

where U are the morphisms such that their composition with the natural projection $(O_{\infty D})^n \rightarrow (O_{\infty})^n$ is an isomorphism. By fixing an isomorphism $\mathbb{F}_q^n \rightarrow (O_{\infty})^n$, U can be identified with the vector bundle

$$V(\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (O_D)^n)).$$

Thus, by Ψ , $\mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D)$ is isomorphic to $V(\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (O_D)^n))$

2.2. Zeta correspondences for vector bundles over $\mathbb{P}_1$ with level structures.

In this section we shall construct a zeta correspondence for $\mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D)$. To do so, we take into account the generalized theta function [An3] 6.1.1.

Over $\mathcal{O}_{\mathbb{P}_1}(k\infty)$ one can consider the following ∞D -level structure, $\pi_{\infty D}^k$:

$$\pi_{\infty D}^k := \pi_{\infty}^k \times \pi_D : \mathcal{O}_{\mathbb{P}_1}(k) \rightarrow \mathcal{O}_{\mathbb{P}_1}/\mathcal{O}_{\mathbb{P}_1}(-1) \times \mathcal{O}_{\mathbb{P}_1}/\mathcal{O}_{\mathbb{P}_1}(-D)$$

π_D being induced by the natural epimorphism $\mathcal{O}_{\mathbb{P}_1} \rightarrow \mathcal{O}_{\mathbb{P}_1}/\mathcal{O}_{\mathbb{P}_1}(-D)$ and the natural inclusion $\mathcal{O}_{\mathbb{P}_1} \hookrightarrow \mathcal{O}_{\mathbb{P}_1}(k)$. π_{∞}^k is obtained from the epimorphism

$$\mathcal{O}_{\mathbb{P}_1}(k) \rightarrow \mathcal{O}_{\mathbb{P}_1}(k)/\mathcal{O}_{\mathbb{P}_1}(k-1)$$

and the isomorphism induced by the multiplication by $1/t^{k-1}$

$$\mathcal{O}_{\mathbb{P}_1}(k)/\mathcal{O}_{\mathbb{P}_1}(k-1) \simeq \mathcal{O}_{\mathbb{P}_1}/\mathcal{O}_{\mathbb{P}_1}(-1).$$

In this way if $\lambda_0 + \lambda_1.t + \dots + \lambda_{k-1}.t^{k-1} \in H^0(\mathbb{P}_1, \mathcal{O}_{\mathbb{P}_1}(k))$,

$$H^0(\pi_{\infty D}^k)(\lambda_0 + \lambda_1.t + \dots + \lambda_{k-1}.t^{k-1}) = (\lambda_0 + \lambda_1.t + \dots + \lambda_{k-1}.t^{k-1}, \lambda_{k-1}) \in O_D \times O_{\infty}$$

Definition 2.5. We define the ∞D -zeta correspondence, $Z_n^{\infty D}$, as the subscheme of $\mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D) \times \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D)$ defined by

$[(\bar{M}, \bar{f}_{\infty D}), (M, f_{\infty D})] \in \text{Hom}_{\text{schemes}}(\text{Spec}(R), Z_n^{\infty D})$ if and only if there exists a morphism of modules $T : \bar{M} \rightarrow M(d-1)$ ($d = \deg(D)$), where the diagram :

$$\begin{array}{ccc} \bar{M} & \xrightarrow{T} & M(d-1) \\ & \searrow \bar{f}_{\infty D} & \downarrow f_{\infty D} \otimes \pi_{\infty D}^{d-1} \\ & & (\mathcal{O}_{\infty D})^n \otimes R \end{array}$$

is commutative.

Therefore, $[(\bar{M}, \bar{f}_{\infty D}), (M, f_{\infty D})] \in Z_n^{\infty D}$ if and only if

$$\text{Hom}_{\text{pairs}}((\bar{M}, \bar{f}_{\infty D}), (M(d-1), f_{\infty D} \otimes \pi_{\infty D}^{d-1})) \neq 0.$$

From Lemma 2.8 it can be deduced that this subscheme is closed.

Lemma 2.6. Given level structures $(\bar{M}, \bar{f}_{\infty D}), (M, f_{\infty D}) \in \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D)$ over R , there exists a unique morphism of vector bundles $T : \bar{M} \rightarrow M(d-1)$ with the diagram

$$\begin{array}{ccc} \bar{M} & \xrightarrow{T} & M(d-1) \\ & \searrow \bar{f}_D & \downarrow f_D \otimes \pi_D \\ & & (\mathcal{O}_D)^n \otimes R \end{array}$$

commutative.

Proof. By choosing bases $\{s_1, \dots, s_n\}$ and $\{\bar{s}_1, \dots, \bar{s}_n\}$ for $H^0(\mathbb{P}_1 \times \text{Spec}(R), M)$ and $H^0(\mathbb{P}_1 \times \text{Spec}(R), \bar{M})$ respectively.

By Remark 1, $(M, f_{\infty D}), (\bar{M}, \bar{f}_{\infty D})$ has associated

$$(\Delta_0 + \Delta_1 t + \dots + \Delta_{d-1} t^{d-1}) \times \Delta_{\infty}$$

and

$$(\bar{\Delta}_0 + \bar{\Delta}_1 t + \dots + \bar{\Delta}_{d-1} t^{d-1}) \times \bar{\Delta}_{\infty},$$

respectively.

Since

$$H^0(\mathbb{P}_1 \times \text{Spec}(R), M(d-1)) = \bigoplus_{i=0}^{d-1} H^0(\mathbb{P}_1 \times \text{Spec}(R), M).t^i$$

by considering the above basis, T is given by

$$A_0 + A_1 t + \dots + A_{d-1} t^{d-1},$$

where A_i are $n \times n$ -matrices with entries in R .

These matrices can be obtained from the relation, $\bar{f}_D = f_D.T|_D$, where $T|_D$ is the isomorphism

$$T|_D : \bar{M}/\bar{M}(-D) \rightarrow M(d-1)/M(d-1)(-D).$$

Thus, T is given by

$$A_0 + A_1 t + \dots + A_{d-1} t^{d-1} = (\bar{\Delta}_0 + \bar{\Delta}_1 t + \dots + \bar{\Delta}_{d-1} t^{d-1})^{-1} \cdot (\Delta_0 + \Delta_1 t + \dots + \Delta_{d-1} t^{d-1}).$$

In this Lemma we have considered an isomorphism $O_D \simeq \mathbb{F}_q[t]/p(t)$ and A_i matrices for linear morphisms between $H^0(\mathbb{P}_1 \times \text{Spec}(R), \bar{M})$ and $H^0(\mathbb{P}_1 \times \text{Spec}(R), M)$ expressed in the bases $\{s_1, \dots, s_n\}$ and $\{\bar{s}_1, \dots, \bar{s}_n\}$. $\square$

Lemma 2.7. With the above notation, $T : \bar{M} \rightarrow M(d-1)$ makes the diagram

$$\begin{array}{ccc} \bar{M} & \xrightarrow{T} & M(d-1) \\ & \searrow \bar{f}_{\infty D} & \downarrow f_{\infty D} \otimes \pi_{\infty D}^{d-1} \\ & & (\mathcal{O}_{\infty D})^n \otimes R \end{array}$$

commutative if and only if $\bar{\Delta}_{\infty} = \Delta_{\infty} \cdot A_{d-1}$.

Proof. By following the same notation as in the latter lemma, let us consider $T = A_0 + A_1 t + \dots + A_{d-1} t^{d-1}$. Then, the only condition that T must satisfy for the diagram

$$\begin{array}{ccc} \bar{M} & \xrightarrow{T} & M(d-1) \\ & \searrow \bar{f}_{\infty D} & \downarrow f_{\infty D} \otimes \pi_{\infty D}^{d-1} \\ & & (\mathcal{O}_{\infty D})^n \otimes R \end{array}$$

to be commutative is that

$$\begin{array}{ccc} \bar{M} & \xrightarrow{T} & M(d-1) \\ & \searrow \bar{f}_{\infty} & \downarrow f_{\infty} \otimes \pi_{\infty}^{d-1} \\ & & (\mathcal{O}_{\infty})^n \otimes R \end{array}$$

must be commutative. But from the definition of π_{∞}^{d-1} , if $i < d-1$ then $\pi_{\infty}^{d-1}(t^i) = 0$ and hence the condition that $A_0 + A_1 t + \dots + A_{d-1} t^{d-1}$ must be satisfied to make the above diagram commutative is $\bar{\Delta}_{\infty} = \Delta_{\infty} \cdot A_{d-1}$. $\square$

Remark 3. From Remark 1

$$V(\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (O_{\infty D})^n)) = \text{Spec}(\mathbb{F}_q[\{d_{i,j,r}\}_{i,j,r}, \{w_{i,\infty}\}_{i,\infty}])$$

" $d_{i,j,r}$ " and " $d_{i,j,\infty}$ " being the matrix entries for $\mathcal{D}_r$ and $\mathcal{D}_{\infty}$, respectively, with

$$(\mathcal{D}_0 + \mathcal{D}_1 t + \dots + \mathcal{D}_{d-1} t^{d-1}) \times \mathcal{D}_{\infty}$$

a universal object for $V := V(\text{Hom}_{\mathbb{F}_q}(\mathbb{F}_q^n, (O_{\infty D})^n))$. Let us denote $p_1, p_2 : V \times V \rightarrow V$ the natural projections and $\mathcal{A}_0 + \mathcal{A}_1 t + \dots + \mathcal{A}_{d-1} t^{d-1}$ the matrix composition

$$(p_2^* \mathcal{D}_0 + p_2^* \mathcal{D}_1 t + \dots + p_2^* \mathcal{D}_{d-1} t^{d-1})^{-1} \cdot (p_1^* \mathcal{D}_0 + \dots + p_1^* \mathcal{D}_{d-1} t^{d-1}).$$

We now consider $I_{\infty D}^n$, the subscheme of $V \times V$, given by the relation

$$p_1^* \mathcal{D}_{\infty} = p_2^* \mathcal{D}_{\infty} \cdot \mathcal{A}_{d-1}.$$

Since $I_{\infty D}^n$ is invariant, by the action of $Gl_n \times Gl_n$, we obtain a subscheme

$$I_{\infty D}^n / Gl_n \times Gl_n \subset V / Gl_n \times V / Gl_n.$$

Moreover, by the very definition of $I_{\infty D}^n$ it is given by the zero locus of n^2 -regular functions of

$$\mathbb{F}_q[\{d_{i,j,r}\}_{i,j,r}, \{d_{i,j,\infty}\}_{i,j}] \otimes \mathbb{F}_q[\{d_{i,j,r}\}_{i,j,r}, \{d_{i,\infty}\}_{i,j}]$$

Lemma 2.8. Bearing in mind the morphism Ψ of 2.4, we have

$$(\Psi \times \Psi)^{-1}(I_{\infty D}^n / Gl_n \times Gl_n) = Z_n^{\infty D}$$

and hence $Z_n^{\infty D}$ is the zero locus of n^2 -regular functions of $\mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D) \times \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D)$. Recall that the subscheme $Z_n^{\infty D}$ is defined in 2.5.

Proof. This Lemma is deduced from Lemma 2.7 and Remark 3.

$Z_n^{\infty D}$ is the zero locus of n^2 -regular functions of $\mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D) \times \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D)$ because of Remark 3. Moreover, one can calculate in an explicit way these n^2 -regular functions from the relation

$$0 = (p_1^* \mathcal{D}_{\infty})^{-1} \cdot p_2^* \mathcal{D}_{\infty} \cdot \mathcal{A}_{d-1}$$

□

For rank 1, we have:

Definition 2.9. We define the ∞D -generalized theta divisor, $\Theta_1^{\infty D}$, as the subscheme of $J_{\mathbb{P}_1, 0}^{\infty D}$ defined by: $(L, f_{\infty D}) \in \text{Hom}_{\text{schemes}}(\text{Spec}(R), \Theta_1^{\infty D})$ if and only if there exists a morphism of modules $T : \mathcal{O}_{\mathbb{P}_1} \otimes R \rightarrow L(d-1)$ ($d = \deg(D)$), where the diagram:

$$\begin{array}{ccc} \mathcal{O}_{\mathbb{P}_1} \otimes R & \xrightarrow{T} & L(d-1) \\ \downarrow \pi_D & \swarrow f_{\infty D} \otimes \pi_{\infty D}^{d-1} & \\ \mathcal{O}_{\infty D} \otimes R & & \end{array}$$

is commutative.

Here, $J_{\mathbb{P}_1, 0}^{\infty D}$ denotes the ∞D -generalized Jacobian of line bundles of degree 0, [Se].

Now, we recall the definition of the tensor product and dual level structures for vector bundles.

Remark 4. The tensor product of two ∞D -level structures $(M, f_{\infty D})$ and $(\bar{M}, \bar{f}_{\infty D})$ is the ∞D -level structure

$$(M \otimes_{\mathcal{O}_{\mathbb{P}_1} \otimes R} \bar{M}, f_{\infty D} \otimes \bar{f}_{\infty D})$$

where $f_{\infty D} \otimes \bar{f}_{\infty D}$ denotes the morphism

$$M \otimes_{\mathcal{O}_{\mathbb{P}_1} \otimes R} \bar{M} \rightarrow (\mathcal{O}_{\infty D})^n \otimes_{\mathcal{O}_{\infty D}} (\mathcal{O}_{\infty D})^n \otimes R \simeq (\mathcal{O}_{\infty D})^{n^2} \otimes R.$$

The dual ∞D -level structure $(M^{\vee}, f_{\infty D}^{\vee})$ for $(M, f_{\infty D})$, is given by a level structure, $f_{\infty D}^{\vee}$, for the dual vector bundle $M^{\vee}$; i.e., $f_{\infty D}$ induces an isomorphism

$$M/M(-\infty - D) \simeq (\mathcal{O}_{\infty D})^n \otimes R$$

from which we deduce an isomorphism

$$M^{\vee} \otimes_{\mathcal{O}_{\mathbb{P}_1} \otimes R} \mathcal{O}_{\infty D} \otimes R \rightarrow (\mathcal{O}_{\infty D})^n \otimes R.$$

$f_{\infty D}^{\vee}$ is given by this isomorphism.

Let us now consider the analogous morphism to the morphism [An2] 4.1. but for line bundles with level structures. Notice that the genus considered in the case of ∞D -level structures is $g(\mathbb{P}_1) + \deg(\infty D) - 1 = d$, [Se] pg.86.

Let $m_{\infty D}$ be the morphism

$$m_{\infty D} : J_{\mathbb{P}_1,0}^{\infty D} \times J_{\mathbb{P}_1,0}^{\infty D} \rightarrow J_{\mathbb{P}_1,0}^{\infty D},$$

defined by

$$m_{\infty D}[(L, f_{\infty D}), (\bar{L}, \bar{f}_{\infty D})] = (\bar{L}, \bar{f}_{\infty D}) \otimes (L^\vee, f_{\infty D}^\vee).$$

We have

Lemma 2.10. For $n = 1$, $m_{\infty D}^{-1}(\Theta_1^{\infty D}) = Z_1^{\infty D}$ and $Z_1^{\infty D}$ is a principal Weil divisor on $J_{\mathbb{P}_1,0}^{\infty D} \times J_{\mathbb{P}_1,0}^{\infty D}$.

Proof. This is deduced from the latter definition, from the definition of $Z_n^{\infty D}$, and from Lemma 2.8. $\square$

3. DRINFELD MODULES AND ZETA CORRESPONDENCES

3.1. A-Drinfeld modules, elliptic sheaves and their antiequivalence. In this section we recall the definition of Drinfeld modules, elliptic sheaves, level structures and their antiequivalence [BlSt], [Dr1], [Dr2],[LRSt], [L],[Mu].

Definition 3.1. A Drinfeld module, ϕ , of rank n over R is a ring morphism:

$$\phi : A \rightarrow \text{End}_R((\mathbb{G}_a)_R)$$

with $\phi_a = \sum_{i=0}^{nv_\infty(a)} a_i \cdot \sigma^i$, $a \in A$, v_∞ is the valuation for ∞ , $a_i \in R$, a_m is a unit in R and $(\mathbb{G}_a)_R$ is the additive line group over R , $\text{End}_R((\mathbb{G}_a)_R)$ are its endomorphisms, and σ is the endomorphism on $(\mathbb{G}_a)_R$, $\sigma(\gamma) = \gamma^q$. In this way, $\text{End}_R((\mathbb{G}_a)_R) = R\{\sigma\}$, where $R\{\sigma\}$ is the ring of polynomials in σ , with the twisted rule of multiplication $\sigma \cdot b = b^q \sigma$.

From this definition, one deduces a morphism of rings

$$c : A \rightarrow R\{\sigma\} \xrightarrow{\sigma=0} R$$

called the characteristic morphism of ϕ .

Definition 3.2. An elliptic sheaf of rank n over R , (E_j, i_j, τ) , is a commutative diagram of vector bundles of rank n over $C \times \text{Spec}(R)$, and injective morphisms of modules $\{i_h\}_{h \in \mathbb{N}}$, τ :

$$\begin{array}{ccccccc} E_1 & \xrightarrow{i_1} & E_2 & \xrightarrow{i_2} & \cdots & \xrightarrow{i_{(n-1)}} & E_n & \xrightarrow{i_n} & \cdots \\ \tau \uparrow & & \tau \uparrow & & \tau \uparrow & & \tau \uparrow & & \\ F^\# E_0 & \xrightarrow{F^\# i_0} & F^\# E_1 & \xrightarrow{F^\# i_1} & \cdots & \xrightarrow{F^\# i_n} & F^\# E_n & \xrightarrow{\quad} & \cdots \end{array}$$

satisfying:

a) For any $z \in \text{Spec}(R)$, $\deg((E_h)_z) = ng + h$.

b) For all $i \in \mathbb{Z}$, $E_{i+n} = E_i(1)$.

$$c) E_i + \tau(F^\# E_i) = E_{i+1}.$$

d) $j_*(E_i/E_{i-1})$ is a rank-one free module over R . j is the inclusion $\infty \times \text{Spec}(R) \hookrightarrow C \times \text{Spec}(R)$.

Recall that $F^\#$ denotes $(Id \times F)^*$.

Remark 5. From these properties, it is deduced that $h^0(E_h) = h$ and $h^1(E_h) = 0$, $h \geq 0$ c.f.[Dr2]. In these conditions it is not hard to prove that the " E_{in} " are semistable.

Moreover, it is proved that there exists a basis $\{s, \sigma.s, \dots, \sigma^{n+i-1}.s\}$ for $H^0(C \times \text{Spec}(R), E_i)$ ($i > 0$), with $\sigma.s := \tau(F^\# s)$ and $\sigma^j.s := \tau(F^\# \sigma^{j-1}.s)$.

In [Dr2], the anti-equivalence between rank- n -Drinfeld modules and rank- n -elliptic sheaves is settled in the following way:

Remark 6. Given a rank- n -Drinfeld module over R

$$\phi : A \rightarrow \text{End}_R((\mathbb{G}_a)_R) = R\{\sigma\}$$

$R\{\sigma\}$ has a structure of an $A \otimes R$ -module by means of ϕ (R -module on the left and A -module on the right). Moreover, $R\{\sigma\}$ has a graduation: $\deg(\sum_j^m b_j.\sigma^j) = m$.

Bearing in mind this graduation, the $A \otimes R$ -module $R\{\sigma\}$ has associated a coherent sheaf E_0 over $C \times \text{Spec}(R)$. The " E_j " are obtained by translating the degree over $R\{\sigma\}$. τ is obtained from the multiplication on the left over $R\{\sigma\}$ by σ . It is not hard to prove that E_i are vector bundles of rank n .

Conversely, given an elliptic sheaf (E_j, i_j, τ) over $C \times \text{Spec}(R)$, by regarding the diagram associated with this elliptic sheaf, for each $h \in \mathbb{N}$ one deduces injective morphisms, which we denote by τ^h

$$\tau^h : F^{\#h} E_i \hookrightarrow E_{i+h}$$

and a chain of modules

$$\tau^k(F^{\#h} E_i) \subset \tau^{k-1}(F^{\#h-1} E_{i+1}) \subset \dots \subset F^{\#h-k} E_{h+k}$$

($h \geq k$).

If $< s > = H^0(C \times \text{Spec}(R), E_{1-n})$ by c) is

$$\bigcup_i H^0(C \times \text{Spec}(R), E_i) = R\{\sigma\}.s$$

with $\sigma^r.s = \tau^r((F^\#)^r(s))$, then $a.s = \sum_j^m b_j.\sigma^j.s$, ($a \in A$) and the Drinfeld module

associated with (E_j, i_j, τ) is $\phi_a = \sum_j^m b_j.\sigma^j$.

Since $\text{rank}(E_i) = \text{rank}(F^\# E_{i-1})$ and $\deg(E_i) = \deg(F^\# E_{i-1}) + 1$, $E_i/\tau(F^\# E_{i-1})$ is a coherent sheaf over $C \times \text{Spec}(R)$ such that for each $s \in \text{Spec}(R)$

$$E_i/\tau(F^\# E_{i-1}) \otimes_R k(s)$$

is concentrated on $p_s \in \text{Spec}(A)$ and hence one obtains a morphism $c^* : \text{Spec}(R) \rightarrow \text{Spec}(A)$. This morphism is the characteristic morphism associated with ϕ . We say that (E_j, i_j, τ) has the generic characteristic if the characteristic morphism $c : A \rightarrow R$ is injective.

Remark 7. Let (E_j, i_j, τ) be an elliptic sheaf defined over a field K and with generic characteristic $c : A \hookrightarrow K$. One can see that $F^{\#h} E_{i+1} / \tau(F^{\#h+1} E_i)$ is concentrated on $y_h \in \text{Spec}(A) \times \text{Spec}(K)$, with $y_0 := (Id \times F^h)(y_h)$ the characteristic of (E_j, i_j, τ) and $E_n / \tau^n(F^{\#n} E_0)$ is concentrated on $y_0, \dots, y_{n-1} \in \text{Spec}(A) \times \text{Spec}(K)$. Moreover, by considering the points " y_i " as morphisms

$$y_i : \text{Spec}(K) \rightarrow \text{Spec}(A) \times \text{Spec}(K)$$

they are given by the morphism $c^{q^i} : A \hookrightarrow K$ on the first entry and the identity morphism on the second entry. Therefore, $y_i \neq y_j$ ($i \neq j$) because $c^{q^i} \neq c^{q^j}$ since c is injective.

Proposition 3.3. Let consider us two elliptic sheaves with generic characteristic (E_j, i_j, τ) and $(\bar{E}_j, \bar{i}_j, \bar{\tau})$ defined over a field K . If $T : E_0 \rightarrow \bar{E}_r$ is a morphism of vector bundles with the diagram

$$\begin{array}{ccc} E_0 & \xrightarrow{T} & \bar{E}_r \\ \tau^n \uparrow & & \bar{\tau}^n \uparrow \\ F^{\#n} E_{-n} & \xrightarrow{F^{\#n} T(-1)} & F^{\#n} \bar{E}_{r-n} \end{array}$$

commutative, there exists a maximum $i \in \mathbb{N}$ such that $T(E_0) \subseteq \bar{\tau}^i(F^{\#i} \bar{E}_{r-i}) \subset \bar{E}_r$. In this case, $(Id \times F^i)y_0 = z_0$, where y_0 and z_0 denote the characteristic of (E_j, i_j, τ) and $(\bar{E}_j, \bar{i}_j, \bar{\tau})$ respectively. We denote by $T(-1)$ the morphism induced by T over $E_{-n} := E_0(-1)$.

Proof. If i is the maximum $i \in \mathbb{N}$ such that $T(E_0) \subseteq \bar{\tau}^i(F^{\#i} \bar{E}_{r-i})$, then we have a non-trivial morphism

$$T : E_0 / F^{\#n} E_{-n} \rightarrow \bar{\tau}^i(F^{\#i} \bar{E}_{r-i}) / \bar{\tau}^{i+1}(F^{\#i+1} \bar{E}_{r-i-1})$$

where $\bar{\tau}^i(F^{\#i} \bar{E}_{r-i}) / \bar{\tau}^{i+1}(F^{\#i+1} \bar{E}_{r-i-1})$ is concentrated on z_i and $E_0 / F^{\#n} E_{-n}$ is concentrated on $y_0, \dots, y_{n-1}$, $(Id \times F^i)z_i = z_0$ and $(Id \times F^j)y_j = y_0$. We thus have $(Id \times F^j)z_0 = (Id \times F^i)y_0$. To conclude it suffices to prove that $j = 0$.

We shall assume that $j \geq 1$. However, in doing so we shall arrive at a contradiction. It is not difficult to see that each case is reduced to studying $j = 1$ and $i = 0$. In this case, $z_0 = y_1$ because $(Id \times F)z_0 = y_0 = (Id \times F)y_1$. Thus,

$$T|_{F^{\#} E_{-1}} : F^{\#} E_{-1} \rightarrow \bar{E}_r / \bar{\tau}(F^{\#} \bar{E}_{r-1})$$

gives an isomorphism

$$F^{\#} E_{-1} / \tau(F^{\#2} E_{-2}) \simeq \bar{E}_r / \bar{\tau}(F^{\#} \bar{E}_{r-1})$$

(recall that $F^\# E_{-1}/\tau(F^{\#^2} E_{-2})$ is concentrated on y_1). We therefore have a commutative diagram

$$\begin{array}{ccc}
 E_0 & \xrightarrow{T} & \bar{E}_r \\
 \uparrow \tau & \nearrow F^\# T_{-1} & \uparrow \bar{\tau} \\
 F^\# E_{-1} & & F^\# E_{r-1} \\
 \uparrow \tau & \nearrow & \\
 F^{\#^2} E_{-2} & &
 \end{array}$$

where $F^\# T_{-1}$ is the restriction of T to $F^\# E_{-1}$. Since $z_0 = y_1$, we have $z_{n-1} = y_n$ because $(Id \times F^{n-1})z_{n-1} = z_0 = y_1 = (Id \times F^{n-1})y_n$ and z_{n-1} and y_n are given by injective morphisms $A \hookrightarrow K$. Thus, in the same way as before we also have a commutative diagram

$$\begin{array}{ccc}
 & & F^{\#^{n-1}} \bar{E}_{r+1-n} \\
 & \nearrow F^{\#^n} T_{-n} & \uparrow \bar{\tau} \\
 F^{\#^n} E_{-n} & \xrightarrow{F^{\#^n} T(-1)} & F^{\#^n} \bar{E}_{r-n} \\
 \uparrow \tau & \nearrow & \\
 F^{\#^{n+1}} E_{-n-1} & &
 \end{array}$$

$F^{\#^n} T_{-n}$ being the restriction of T to $F^{\#^n} E_{-n}$. Thus, this morphism gives an isomorphism

$$F^{\#^n} E_{-n}/\tau(F^{\#^{n+1}} E_{-n-1}) \simeq F^{\#^{n-1}} \bar{E}_{r-n+1}/\bar{\tau}(F^{\#^n} \bar{E}_{r-n}).$$

However, this contradicts the inclusion

$$F^{\#^n} T(-1)(F^{\#^n} E_{-n}) \subseteq F^{\#^n} \bar{E}_{r-n}.$$

□

3.2. A -Drinfeld modules and elliptic sheaves with level structures. Let us now briefly recall the definitions of level structures for elliptic sheaves and Drinfeld modules: (E_j, i_j, τ) and ϕ respectively.

Let E_I be the subgroup scheme of I division points of $(\mathbb{G}_a)_R$ as an A -module (via ϕ). $(\widetilde{I^{-1}/A})^n$ will denote the constant sheaf of stalk $(I^{-1}/A)^n$. I is a proper ideal in $\text{Spec}(A)$.

Definition 3.4. An I -level structure for ϕ is a pair (ϕ, ι_I) . ι_I is an isomorphism of A -modules

$$\iota_I : E_I(R) \rightarrow (\widetilde{I^{-1}/A})^n(R).$$

Definition 3.5. A D -level structure for the elliptic sheaf (E_j, i_j, τ) , is a D -level structure, f_D^j , for each vector bundle E_j compatible with the morphisms $\{i_j, \tau\}$. i.e., $f_{rx}^D \cdot i_j = f_D^j$ and $f_D^{j+1} \cdot \tau = (Id \times F)^* f_D^j$.

In [An1], the anti-equivalence between Drinfeld modules and elliptic sheaves with level structures is established. There exists an affine scheme, $\mathcal{E}_n^D = \text{Spec}(\mathfrak{R}_n^D)$, that represents rank- n Elliptic sheaves with D -level structures (equivalently, rank- n Drinfeld modules with I -level structures).

One can calculate in an explicit way the sections $\sigma^i.s$ (Remark 5) via level structures. These computations are done in [An1](Theorem 5), [Al] (Remark 3.1).

Let I be the ideal associated with the effective divisor $D \subset \text{Spec}(A)$, $D = r_1.x_1 + \dots + r_l.x_l$, x_i being the point in $\text{Spec}(A)$ associated with the maximal ideal m_{x_i} and t_{x_i} a local uniformizer for m_{x_i} .

Remark 8. if (Φ, ι_I) is a universal Drinfeld module with an I -level structure over $\mathfrak{R}_n^D$. Therefore, ι_I is an isomorphism

$$\iota_I : E_I(\mathfrak{R}_n^D) \rightarrow \left(\prod_{i \in \{1, \dots, l\}} t_{x_i}^{-r_i} \mathbb{F}_q[[t_{x_i}]] / \mathbb{F}_q[[t_{x_i}]] \right)^n$$

with $\iota_I(\alpha_{jh}^{x_i}) = (0, \dots, \overset{j}{v_j}, \dots, 0)$ and

$$v_j = (0, \dots, \overset{i}{t}^{-1-h}, \dots, 0) \in \prod_{i \in \{1, \dots, l\}} t_{x_i}^{-r_i} \mathbb{F}_q[[t_{x_i}]] / \mathbb{F}_q[[t_{x_i}]].$$

We denote by $(\mathbb{E}_j, i_j, \tau, f_D^j)$ the corresponding universal elliptic sheaf of rank n with a D -level structure associated with (Φ, ι_I) . Thus, the value of the sections $\sigma^k s \in H^0(C \otimes \mathfrak{R}_n^D, \mathbb{E}_m)$ via the level structure $f_D^m : \mathbb{E}_m \rightarrow \mathcal{O}_D^n \otimes \mathfrak{R}_n^D$ is:

$$H^0(f_D^m)(\sigma^k s) = (s_1^k, \dots, s_n^k) \in (\mathcal{O}_D \otimes \mathfrak{R}_n^D)^n$$

By using the isomorphism

$$(\mathcal{O}_D \otimes \mathfrak{R}_n^D)^n = \left(\prod_{i \in \{1, \dots, l\}} \mathfrak{R}_n^D[[t_{x_i}]] / t_{x_i}^{r_i} \mathfrak{R}_n^D[[t_{x_i}]] \right)^n$$

we have:

$$s_j^k = \left(\sum_{h=0}^{r_1-1} (\alpha_{jh}^{x_1})^{q^k} t_{x_1}^h, \dots, \sum_{h=0}^{r_l-1} (\alpha_{jh}^{x_r})^{q^k} t_{x_r}^h \right).$$

Definition 3.6. A ∞ -level structure for a rank- n elliptic sheaf (E_j, i_j, τ) over R is a ∞ -level structure (E_0, f_∞) such that the diagram

$$\begin{array}{ccc} F^{\#n} E_0 & \xrightarrow{\tau^n} & E_0(1) \\ & \searrow F^{\#n} f_\infty & \downarrow f_\infty \otimes \pi_\infty \\ & & (\mathcal{O}_\infty)^n \otimes R \end{array}$$

is commutative. π_∞ is defined in 2.2

To give a ∞D -level structure for an elliptic sheaf is to give an ∞ -level structure together with a D -level structure for the elliptic sheaf.

The idea of considering ∞ -level structures in this setting has been suggested to me by G.W. Anderson.

As for D -level structures, there exists an affine scheme, $\mathcal{E}_n^{\infty D} = \text{Spec}(\mathfrak{R}_n^{\infty D})$, that represents rank- n Elliptic sheaves with ∞D -level structures.

Remark 9. Let $(\mathbb{E}_j, i_j, \tau)$ be a universal rank- n elliptic sheaf with an ∞D -level structure $f_{\infty D} = f_{\infty} \times f_D$ over $\text{Spec}(\mathfrak{R}_n^{\infty D})$. By choosing the basis

$$\{s, \sigma.s, \dots, \sigma^{n-1}.s\}$$

in $H^0(C \otimes \mathfrak{R}_n^{\infty D}, \mathbb{E}_0)$, Remark 1 associates a matrix with $(\mathbb{E}_0, f_{\infty D})$. By using the results and notation of the latter Remark, this matrix is

$$(A_{x_1,0} + A_{x_1,1}.t_{x_1} + \dots + A_{x_1,r_1-1}.t_{x_1}^{r_1-1}) \times \dots \times (A_{x_l,0} + A_{x_l,1}.t_{x_l} + \dots + A_{x_l,r_l-1}.t_{x_l}^{r_l-1}) \times A_{\infty},$$

where $A_{x_i,h}$ is

$$\begin{pmatrix} \alpha_{1,h}^{x_i} & (\alpha_{1,h}^{x_i})^q & \dots & (\alpha_{1,h}^{x_i})^{q^{n-1}} \\ \alpha_{2,h}^{x_i} & (\alpha_{2,h}^{x_i})^q & \dots & (\alpha_{2,h}^{x_i})^{q^{n-1}} \\ \dots & \dots & \dots & \dots \\ \alpha_{n,h}^{x_i} & (\alpha_{n,h}^{x_i})^q & \dots & (\alpha_{n,h}^{x_i})^{q^{n-1}} \end{pmatrix}.$$

A_{∞} is a $n \times n$ -matrix with coefficients in $\mathfrak{R}_n^{\infty D}$ that we shall calculate in the case $C = \mathbb{P}_1$ in the following subsection.

3.3. $\mathbb{F}_q[t]$ -Drinfeld modules and zeta correspondences. In the rest of this section $C = \mathbb{P}_1$. We denote by $\mathcal{E}_n^{\infty D}$ the moduli (scheme) of rank- n elliptic sheaves with ∞D -level structures for $A = \mathbb{F}_q[t]$. We denote an object of this moduli by $(E_j, i_j, \tau, f_{\infty D})$.

In a direct way, one can obtain a morphism

$$\Phi : \mathcal{E}_n^{\infty D} \rightarrow \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D)$$

given by $\Phi(E_j, i_j, \tau, f_{\infty D}) = (E_0, f_{\infty} \times f_D^0)$. Recall that by Remark 5, E_0 are semistable.

The next three subsections are devoted to studying the zeta correspondences

$$(\Phi \times \Phi)^{-1}(Z_n^{\infty D})$$

in $\mathcal{E}_n^{\infty D} \times \mathcal{E}_n^{\infty D}$, where $Z_n^{\infty D}$ is defined as in Definition 2.5.

$\Phi \times \Phi$ is the analogous morphism to that defined in [An2] 4.1.

Lemma 3.7. If

$$[(E_j, i_j, \tau, f_{\infty D}), (\bar{E}_j, \bar{i}_j, \bar{\tau}, \bar{f}_{\infty D})] \in (\Phi \times \Phi)^{-1}(Z_n^{\infty D}),$$

then there exists $h \in \mathbb{N}$ with $h \leq n(d-1)$ such that $(Id \times F^h)y_0 = z_0$, y_0 and z_0 being the characteristics of (E_j, i_j, τ) and $(\bar{E}_j, \bar{i}_j, \bar{\tau})$ respectively.

Proof. We assume the elliptic sheaves defined over a field R . By Definition 2.5, there exists a diagram (not necessarily commutative)

$$\begin{array}{ccc}
 E_0 & \xrightarrow{T} & \bar{E}_{n(d-1)} \\
 \tau^n \uparrow & \searrow & \swarrow \uparrow \bar{\tau}^n \\
 & (\mathcal{O}_{\infty D})^{\otimes} R & \\
 & \swarrow & \searrow \\
 F^{\#n} E_{-n} & \xrightarrow{F^{\#n} T(-1)} & F^{\#n} \bar{E}_{n(d-2)}
 \end{array}$$

(oblique arrows are given by level structures). This diagram is commutative after tensoring by $\mathcal{O}_{\infty D}$ and we therefore have that the morphism of vector bundles

$$T \cdot \tau^n - \bar{\tau}^n \cdot F^{\#n} T(-1) : F^{\#n} E_{-n} \rightarrow \bar{E}_{n(d-1)}$$

takes values in $\bar{E}_{n(d-1)}(-\infty - D)$. However, we are in $\mathbb{P}_1$ and $\bar{E}_{n(d-1)}$ and $F^{\#n} E_{-n}$ are semistable. Therefore,

$$\bar{E}_{n(d-1)} \otimes_{\mathbb{P}_1 \otimes R} (F^{\#n} E_{-n})^{\vee}(-\infty - D)$$

is also semistable of degree $-n$. In this way,

$$H^0(\mathbb{P}_1 \otimes R, \bar{E}_{n(d-1)} \otimes_{\mathbb{P}_1 \otimes R} (F^{\#n} E_{-n})^{\vee}(-\infty - D)) = 0.$$

Hence $T \cdot \tau^n - \bar{\tau}^n \cdot F^{\#n} T(-1) = 0$. We therefore conclude because we are in the conditions of Proposition 3.3. $\square$

Let K be the $\mathbb{F}_q(t)$ -algebra, $(t \rightarrow x)$, $K := \mathbb{F}_q(x)[a_1, \dots, a_{n-1}]$. Let ϕ be the rank- n Drinfeld module with generic characteristic, defined over K :

$$\phi_t = \sigma^n + a_{n-1} \cdot \sigma^{n-1} + \dots + a_1 \cdot \sigma + x.$$

Let (E_j, i_j, τ) be an elliptic sheaf associated with ϕ and $s \in H^0(\mathbb{P}_1 \otimes K, E_{1-n})$, with $t \cdot s = x \cdot s + a_1 \cdot \sigma \cdot s + \dots + a_{n-1} \cdot \sigma^{n-1} \cdot s + \sigma^n \cdot s$.

Bearing in mind Definition 3.6, we study when an ∞ -level structure f_{∞} for E_0 is an ∞ -level structure for the elliptic sheaf (E_j, i_j, τ) .

Since $H^0(\mathbb{P}_1 \otimes K, E_0(1)) = H^0(\mathbb{P}_1 \otimes K, E_0) \oplus t \cdot H^0(\mathbb{P}_1 \otimes K, E_0)$ if we consider the bases $\{s, \sigma \cdot s, \dots, \sigma^{n-1} \cdot s\}$ and $\{F^{\#n}(s), \sigma \cdot F^{\#n}(s), \dots, \sigma^{n-1} \cdot F^{\#n}(s)\}$ for $H^0(\mathbb{P}_1 \otimes K, E_0)$ and $H^0(\mathbb{P}_1 \otimes K, F^{\#} E_0)$ respectively, the morphism $\tau^n : F^{\#n} E_0 \rightarrow E_0(1)$ is given in these basis by $A \cdot t + B$, A, B being matrices with entries in K .

Lemma 3.8. We have that

$$A \cdot t + B = F^{\#n-1} C \cdot F^{\#n-2} C \dots F^{\#} C \cdot C$$

with

$$C = \begin{pmatrix} 0 & 0 & \dots & 0 & t - x \\ 1 & 0 & \dots & 0 & -a_{n-1} \\ 0 & 1 & \dots & 0 & -a_{n-2} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & -a_1 \end{pmatrix}$$

Proof. Bearing in mind Remark 6, τ^n is given by

$$F^{*n}K\{\sigma\} \xrightarrow{\sigma^n \cdot \sigma} K\{\sigma\},$$

where $\sigma \cdot$ is the multiplication on the left over $K\{\sigma\}$. In this setting, the bases $\{s, \sigma.s, \dots, \sigma^{n-1}.s\}$ and $\{F^{\#n}(s), F^{\#n}(\sigma.s), \dots, F^{\#n}(\sigma^{n-1}.s)\}$ are $\{1, \sigma, \dots, \sigma^{n-1}\}$ and $\{F^{\#n}(1), F^{\#n}(\sigma), \dots, F^{\#n}(\sigma^{n-1})\}$ respectively. We conclude bearing in mind that $t.s = x.s + a_1.\sigma.s + \dots + a_{n-1}.\sigma^{n-1}.s + \sigma^n.s$. $\square$

Now we shall calculate A_∞ (Remark 9)

Lemma 3.9. Let (E_j, i_j, τ) be a rank- n elliptic sheaf, f_∞ an ∞ -level structure over E_0 , and $(\nu_{ij})_{i,j}$ the matrix associated with

$$f_\infty : H^0(\mathbb{P}_1 \otimes K, E_0) \rightarrow K^{\oplus n}$$

for the basis $\{s, \sigma.s, \dots, \sigma^{n-1}.s\}$ and the standard basis in K^n . Then (E_0, f_∞) is an ∞ -level structure for the elliptic sheaf (E_j, i_j, τ) if and only if $(\nu_{ij}^q)_{i,j} = (\nu_{ij})_{i,j} \cdot A$, where A is defined as in the above Lemma.

Proof. Because of the definition of ∞ -level structures for elliptic sheaves, the diagram

$$\begin{array}{ccc} F^{\#n}E_0 & \xrightarrow{\tau^n} & E_0(1) \\ & \searrow F^{\#n}f_\infty & \downarrow f_\infty \otimes \pi_\infty \\ & & (\mathcal{O}_\infty)^n \otimes R \end{array}$$

must be commutative.

Bearing in mind 1) of Proposition 2.3 and the above Lemma, we conclude by taking the bases $\{F^{\#n}(s), F^{\#n}(\sigma.s), \dots, F^{\#n}(\sigma^{n-1}.s)\}$ in $H^0(\mathbb{P}_1 \otimes K, F^{\#n}E_0)$ and

$$\{s, \sigma.s, \dots, \sigma^{n-1}.s\} \cup t.\{s, \sigma.s, \dots, \sigma^{n-1}.s\}$$

for $H^0(\mathbb{P}_1 \otimes K, E_0(1))$. $\square$

A_∞ (Remark 9) is $(\nu_{ij})_{i,j}$.

We shall now consider $\bar{K}_n^{\infty D} := K[\{\alpha_{jh}^{x_i}\}_{i,j,h}, \{\nu_{ij}\}_{i,j}]$ as a subring of an algebraic closed field containing K , where the " $\alpha_{jh}^{x_i}$ " are defined as in Remark 8 and " ν_{ij} " are solutions for the equation $(\nu_{ij}^q)_{i,j} = (\nu_{ij})_{i,j} \cdot A$.

It is not hard to prove that there exists a morphism $\text{Spec}(\bar{K}_n^{\infty D}) \rightarrow \mathcal{E}_n^{\infty D}$.

Theorem 3.10. If we denote by β the morphism composition

$$\text{Spec}(\bar{K}_{n,\infty D}) \times \text{Spec}(\bar{K}_{n,\infty D}) \longrightarrow \mathcal{E}_n^{\infty D} \times \mathcal{E}_n^{\infty D} \xrightarrow{\Phi \times \Phi} \mathcal{M}_{\mathbb{P}^1}^{ss}(n, 0, \infty D) \times \mathcal{M}_{\mathbb{P}^1}^{ss}(n, 0, \infty D),$$

then

$$\beta^{-1}(Z_n^{\infty D})$$

is the zero locus of the n^2 -functions deduced from an equation

$$\bar{\Gamma}_\infty \cdot \bar{\Gamma}_{d-1} - Id_n = 0,$$

$\bar{\Gamma}_{d-1}, \bar{\Gamma}_\infty$ being $n \times n$ -matrices with entries in $\bar{K}_n^{\infty D} \otimes \bar{K}_n^{\infty D}$.

Proof. This Theorem is a direct consequence of Lemma 2.8 and one can make an explicit calculation of $\bar{\Gamma}_{d-1}, \bar{\Gamma}_\infty$ bearing in mind Remark 9 for parameters $t_{x_i} = t - \alpha_i$, Lemma 3.9, and the Chinese remainder theorem: i.e., there exists a ring isomorphism

$$\delta : \prod_{i \in \{1, \dots, l\}} \mathbb{F}_q[t]/(t - \alpha_i)^{r_i} \rightarrow O_D := \mathbb{F}_q[t]/(p(t))$$

given by

$$\delta(h_1(t), \dots, h_l(t)) = p_1(t - \alpha_1).h_1(t).p(t)/(t - \alpha_1)^{r_1} + \dots + p_l(t - \alpha_l).h_l(t).p(t)/(t - \alpha_l)^{r_l}$$

with $\deg(p_i(t - \alpha_i)) < r_i$ and

$$1/p(t) = \sum_{i \in \{1, \dots, l\}} p_i(t - \alpha_i)/(t - \alpha_i)^{r_i}.$$

Recall that $D = (p(t))_0$, with $p(t) = \prod_{i \in \{1, \dots, l\}} (t - \alpha_i)^{r_i}$.

□

3.4. Some explicit irreducible components for zeta correspondences. The finite group $Gl_{\mathcal{O}_{\infty D}}(\mathcal{O}_{\infty D}^n) = Gl_n(\mathbb{F}_q)_\infty \times Gl_{\mathcal{O}_D}(\mathcal{O}_D^n)$ acts on $\tilde{K}_n^{\infty D}$ by acting on the level structures. By considering the restrictions over $\tilde{K}_n^{\infty D}$

$$a_1 = q_1(x), \dots, a_{n-1} = q_{n-1}(x)$$

with $q_1(x), \dots, q_{n-1}(x) \in \mathbb{F}_q[x]$, if we denote

$$\tilde{K}_n^{\infty D} := \tilde{K}_n^{\infty D} / (a_1 - q_1(x), \dots, a_{n-1} - q_{n-1}(x))$$

we have

$$Spec(\tilde{K}_n^{\infty D}) / Gl_n(\mathbb{F}_q)_\infty \times Gl_{\mathcal{O}_D}(\mathcal{O}_D^n) = Spec(\mathbb{F}_q(x)).$$

Lemma 3.11. Let $\tilde{\beta}$ be the restriction of β to $Spec(\tilde{K}_n^{\infty D}) \times Spec(\tilde{K}_n^{\infty D})$:

$$\tilde{\beta} : Spec(\tilde{K}_n^{\infty D}) \times Spec(\tilde{K}_n^{\infty D}) \longrightarrow \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D) \times \mathcal{M}_{\mathbb{P}_1}^{ss}(n, 0, \infty D).$$

Then,

$$(\tilde{\beta})^{-1}(Z_n^{\infty D}) = \bigcup_{\substack{\text{For certain } g \in Gl_{\mathcal{O}_{\infty D}}(\mathcal{O}_{\infty D}^n) \\ \text{and } 0 \leq i \leq n(d-1)}} \Gamma_{g.F^i}.$$

F is the Frobenius morphism and $\Gamma_{g.F^i}$ is the graph for the morphism $g.F^i$. g is considered as an automorphism over $\tilde{K}_n^{\infty D}$.

Proof. Let $(E_j, i_j, \tau, f_{\infty D})$ and $(\bar{E}_j, \bar{i}_j, \bar{\tau}, \bar{f}_{\infty D})$ be two rank- n elliptic sheaves with an ∞D -level structure defined over $\tilde{K}_n^{\infty D}$. For Proposition 3.3, if

$$[(E_0, f_{\infty D}), (\bar{E}_0, \bar{f}_{\infty D})] \in (\tilde{\beta})^{-1}(Z_n^{\infty D}),$$

then F^j (Characteristic($E_j, i_j, \tau, f_{\infty D}$)) = Characteristic($\bar{E}_j, \bar{i}_j, \bar{\tau}, \bar{f}_{\infty D}$) for some $j \in \mathbb{N}$. Thus, for some $g \in Gl_{\mathcal{O}_{\infty D}}(\mathcal{O}_{\infty D}^n)$ is

$$g.F^j(E_j, i_j, \tau, f_{\infty D}) = (\bar{E}_j, \bar{i}_j, \bar{\tau}, \bar{f}_{\infty D})$$

because

$$Spec(\tilde{K}_n^{\infty D}) \rightarrow Spec(\tilde{K}_n^{\infty D}) / Gl_{\mathcal{O}_{\infty D}}(\mathcal{O}_{\infty D}^n) = Spec(\mathbb{F}_q(x))$$

is the characteristic morphism.

□

Lemma 3.12. Let $s(t)$ be a monic polynomial with $d - 1 - i = \deg(s(t)) \leq d - 1$. We denote by $h_{s(t)}$ the central matrix $Id_n \times \text{diag}(s(t), \dots, s(t)) \in Gl_n(\mathbb{F}_q)_\infty \times Gl_{\mathcal{O}_D}(\mathcal{O}_D^n)$. Then,

$$\Gamma_{h_{s(t)} \cdot F^{ni}} \subset (\tilde{\beta})^{-1}(Z_n^{\infty D}).$$

Moreover, if

$$\Gamma_{g \cdot F^{n(d-1)}} \subset (\tilde{\beta})^{-1}(Z_n^{\infty D}),$$

then $g = Id$.

Proof. It suffices to find $T : F^{\#ni}(E_0) \rightarrow E_0(d-1)$ such that the diagram

$$\begin{array}{ccc} F^{\#ni}(E_0) & \xrightarrow{T} & E_0(d-1) \\ & \searrow h_{s(t)} \cdot F^{\#ni} f_{\infty D} & \downarrow f_{\infty D} \otimes \pi_{\infty D}^{d-1} \\ & & (\mathcal{O}_{\infty D})^n \otimes \tilde{K}_n^{\infty D} \end{array}$$

is commutative. However, since

$$\begin{array}{ccc} F^{\#ni}(E_0) & \xrightarrow{\tau^{ni}} & E_0(i) \\ & \searrow F^{\#ni} f_{\infty D} \otimes \pi_{\infty D}^i & \downarrow f_{\infty D} \otimes \pi_{\infty D}^{d-1} \\ & & (\mathcal{O}_{\infty D})^n \otimes \tilde{K}_n^{\infty D} \end{array}$$

is commutative because $f_{\infty D}$ is an ∞D -level structure for the elliptic sheaf (E_j, i_j, τ) , it suffices to find $T' : E_0(i) \rightarrow E_0(d-1)$ with

$$\begin{array}{ccc} E_0(i) & \xrightarrow{T'} & E_0(d-1) \\ & \searrow h_{s(t)} \cdot (f_{\infty D}) \otimes \pi_{\infty D}^i & \downarrow f_{\infty D} \otimes \pi_{\infty D}^{d-1} \\ & & (\mathcal{O}_{\infty D})^n \otimes \tilde{K}_n^{\infty D} \end{array}$$

commutative. We shall take T' defined as the homothety by $s(t)$. It is clear that it is defined from $E_0(i)$ to $E_0(d-1)$, because $\deg(s(t)) = d - 1 - i$. Moreover, the above diagram is commutative because

$$\begin{array}{ccc} E_0(i) & \xrightarrow{T'} & E_0(d-1) \\ & \searrow h_{s(t)} \cdot (f_D) \otimes \pi_D & \downarrow f_D \otimes \pi_D \\ & & (\mathcal{O}_D)^n \otimes \tilde{K}_n^{\infty D} \end{array}$$

is commutative by the definition of T' . And

$$\begin{array}{ccc} E_0(i) & \xrightarrow{T'} & E_0(d-1) \\ & \searrow h_{s(t)} \cdot (f_{\infty}) \otimes \pi_{\infty}^i & \downarrow f_{\infty} \otimes \pi_{\infty}^{d-1} \\ & & (\mathcal{O}_{\infty})^n \otimes \tilde{K}_n^{\infty D} \end{array}$$

is commutative because $s(t)$ is monic.

The second assertion of the Lemma is deduced because T takes values in

$$\tau^{n(d-1)}(E_0^{\sigma^{n(d-1)}}) \subset E_{n(d-1)}$$

because of Proposition 3.3. We thus have $\tau^{n(d-1)} = T$, (up to isomorphisms). In this case, if

$$(f_{\infty D} \otimes \pi_{\infty D}^{d-1}).\tau^{n(d-1)} = g.F^{\#n(d-1)}f_{\infty D},$$

this implies that $g = Id$ because the definition of ∞D -level structure for an elliptic sheaf is

$$(f_{\infty D} \otimes \pi_{\infty D}^{d-1}).\tau^{n(d-1)} = F^{\#n(d-1)}f_{\infty D}.$$

□

From this Lemma we deduce:

Theorem 3.13. (c.f [An3],[AnDP], [Si]) For $n = 1$

$$(\tilde{\beta}_1)^{-1}(\Theta_1^{\infty D}) = \bigcup_{\substack{s(t) \in (\mathbb{F}_q[t]/p(t))^\times \\ s(t) \text{ monic with } \deg(s(t))=i \\ 0 \leq i \leq d-1}} \Gamma_{s(t).F^i}$$

Here, $\tilde{\beta}_1$ denotes $m_{\infty D}.\tilde{\beta}$ and we follow the notation of Lemma 2.10.

We can now obtain the same result as in [An3] merely by considering all possibilities over the ∞ -level structures: From Lemma 3.9 for rank 1, two ∞ -level structures for a rank-1 elliptic sheaf differ in a $a \in \mathbb{F}_q^\times$. Thus,

$$\bigcup_{a \in \mathbb{F}_q^\times} (\tilde{\beta}_1)^{-1}(\Theta_{1,a \times 1}^{\infty D}) = \bigcup_{\substack{s(t) \in (\mathbb{F}_q[t]/p(t))^\times \\ 0 \leq i \leq d-1, \deg(s(t))=i}} \Gamma_{s(t).F^i}$$

Here, $a \times 1 \in \mathbb{F}_q^\times \times \mathcal{O}_D^\times$, $a \times 1$ acts on $\mathcal{M}_{\mathbb{P}^1}^{ss}(1, 0, \infty D)$ (this scheme is the generalized Jacobian $J_{\mathbb{P}^1}^{\infty D}$), and $\Theta_{1,a \times 1}^{\infty D}$ denotes the transformation of $\Theta_1^{\infty D}$ by $a \times 1$.

3.5. Examples. As above, C will be $\mathbb{P}_1$.

Example. 1 ([An4],[C]) We shall study $n = 1$, $p(t) = t(t-1)$. We thus denote D by $0+1$, and $\tilde{K}_1^{\infty+0+1} = \mathbb{F}_q(x)[\alpha_0, \alpha_1, \nu]$, where

$$\alpha_0^q + \alpha_0.x = 0, \alpha_1^q + \alpha_1.(x-1) = 0,$$

and $\nu \in \mathbb{F}_q^\times$. We can set $\nu = 1$. With the variable changes $u = \alpha_0$ and $v = \alpha_1$, we have

$$\mathbb{F}_q(x)[\alpha_0, \alpha_1, \nu] = \mathbb{F}_q(u)[v, 1/v]/u^{q-1} - v^{q-1} + 1$$

Now bearing in mind the Chinese remainder theorem, there exists a ring isomorphism

$$\delta : \mathbb{F}_q[t]/(t) \times \mathbb{F}_q[t]/(t-1) = \mathbb{F}_q[t]/(t(t-1)),$$

given by

$$\delta(c, d) = ct + d(1-t).$$

In this way, $ct + d(1-t)$ is monic $\in (\mathbb{F}_q[t]/(t(t-1)))^\times$ if and only if $c-d=1$ and $c \neq 0, 1$.

For Theorem 3.13,

$$(\tilde{\beta}_1)^{-1}(\Theta_1^{\infty+0+1}) = \bigcup_{\substack{c \in \mathbb{F}_q^\times \\ c \neq 0, 1}} \Gamma_{[c, c-1]} \cup \Gamma_F,$$

$\Gamma_{[c,c-1]}$ being the graph of the morphism in $\tilde{K}_1^{\infty+0+1}$ defined by $u \rightarrow c.u$, $v \rightarrow (c-1).v$ and

$$\tilde{\beta}_1 : \text{Spec}(\tilde{K}_1^{\infty+0+1}) \times \text{Spec}(\tilde{K}_1^{\infty+0+1}) \longrightarrow \mathcal{M}_{\mathbb{P}_1}^{ss}(1, 0, \infty + 0 + 1) = J_{\infty+0+1}^{\mathbb{P}_1},$$

defined in 3.13. By Theorem 3.10 and by the explicit calculations in Remark 9 and Lemma 3.9, $\tilde{\beta}_1$ has associated the 1×1 -”matrix”

$$(1 \otimes u \times 1 \otimes v \times 1 \otimes 1)^{-1} \cdot (u \otimes 1 \times v \otimes 1 \times 1 \otimes 1)$$

and

$$(\tilde{\beta}_1)^{-1}(\Theta_1^{\infty+0+1}) = \left(\frac{u \otimes 1}{1 \otimes u} - \frac{v \otimes 1}{1 \otimes v} - 1 \otimes 1 \right)_0.$$

If we consider $\nu = a \in \mathbb{F}_q^\times$, we obtain

$$(\tilde{\beta}_1)^{-1}(\Theta_{1,1 \times a}^{\infty+0+1}) = \left(\frac{u \otimes 1}{1 \otimes u} - \frac{v \otimes 1}{1 \otimes v} - a \otimes 1 \right)_0$$

and

$$(\tilde{\beta}_1)^{-1}(\Theta_{\infty+0+1}^{1,1 \times a}) = \bigcup_{\substack{c \in \mathbb{F}_q^\times \\ c \neq 0, a}} \Gamma_{[c,c-a]} \cup \Gamma_F$$

Example. 2 In this example we shall study the last example in a more general way. $n = 1$, $p(t) = \prod_{i \in \{1, \dots, l\}} (t - \alpha_i)$. Then, $D = \alpha_1 + \dots + \alpha_l$ and

$$\tilde{K}_1^{\infty+D} = \mathbb{F}_q(x)[\alpha_1, \dots, \alpha_l, \nu],$$

where $\alpha_i^q + \alpha_i \cdot (x - \alpha_i) = 0$ with $1 \leq i \leq l$ and $\nu \in \mathbb{F}_q^\times$. We can set $\nu = 1$ as in the above example.

Let us consider the decomposition in simple fractions

$$1/p(t) = \sum_{i \in \{1, \dots, l\}} m_i/t - \alpha_i,$$

$m_i \in \mathbb{F}_q^\times$. The Chinese remainder theorem is then settled in the following way: There exists a ring isomorphism

$$\delta : \prod_{i \in \{1, \dots, l\}} \mathbb{F}_q[t]/(t - \alpha_i) \rightarrow \mathbb{F}_q[t]/(p(t))$$

given by

$$\delta(h_1, \dots, h_l) = m_1.h_1.p(t)/t - \alpha_1 + \dots + m_l.h_l.p(t)/t - \alpha_l.$$

In this way, $\delta(h_1, \dots, h_l) \in (\mathbb{F}_q[t]/(p(t)))^\times$ if $h_i \neq 0$ for all i .

For Theorem 3.13 $(\tilde{\beta}_1)^{-1}(\Theta_1^{\infty+\alpha_1+\dots+\alpha_l})$ is

$$\bigcup_{0 \leq i \leq \deg(p(t))} \bigcup_{\substack{(h_1, \dots, h_l) \\ \deg(\delta(h_1, \dots, h_l)) = \deg(p(t)) - i \\ \delta(h_1, \dots, h_l) \text{ monic and } h_1 \dots h_l \neq 0}} \Gamma_{\delta(h_1, \dots, h_l).F^i}$$

$\Gamma_{\delta(h_1, \dots, h_l).F^i}$ is the graph of the morphism in $\tilde{K}_1^{\infty+0+1}$ defined by $\alpha_r \rightarrow h_r \cdot \alpha_r^{q^i}$, for each i .

By Theorem 3.10 and by the explicit calculations in Remark 9 and Lemma 3.9, $\tilde{\beta}_1$ has associated the 1×1 -”matrix”

$$(1 \otimes \alpha_1 \times \dots \times 1 \otimes \alpha_l \times 1 \otimes 1)^{-1} \cdot (\alpha_1 \otimes 1 \times \dots \times \alpha_l \otimes 1 \times 1 \otimes 1)$$

and

$$(\tilde{\beta}_1)^{-1}(\Theta_1^{\infty+\alpha_1+\dots+\alpha_l}) = (m_1 \cdot \frac{\alpha_1 \otimes 1}{1 \otimes \alpha_1} + \dots + m_l \cdot \frac{\alpha_l \otimes 1}{1 \otimes \alpha_l} - 1 \otimes 1)_0.$$

Example. 3 We now consider $n = 2$, $p(t) = t$. Then, $D = 0$ and we follow the notation of Remark 9

$$\tilde{K}_2^{\infty+0} = \mathbb{F}_q(x)[\alpha_{1,0}^0, \alpha_{2,0}^0, \nu_{11}, \nu_{21}, \nu_{12}, \nu_{22}]$$

where

$$(\alpha_{i,0}^0)^{q^2} + q_1(x) \cdot (\alpha_{i,0}^0)^q + \alpha_{i,0}^0 \cdot x = 0$$

for $i = 1, 2$ and $(\nu_{ij})_{ij}$ satisfies $(\nu_{ij}^{q^2})_{ij} = (\nu_{ij})_{ij} \cdot A$, with A defined as in Lemma 3.9

$$At + B = \begin{pmatrix} 0 & t - x^q \\ 1 & -q_1(x)^q \end{pmatrix} \cdot \begin{pmatrix} 0 & t - x \\ 1 & -q_1(x) \end{pmatrix}.$$

Thus,

$$A = \begin{pmatrix} 1 & -q_1(x) \\ 0 & 1 \end{pmatrix},$$

and hence $\nu_{21}^{q^2} = \nu_{21}$, $\nu_{22}^{q^2} = \nu_{22}$ and $\nu_{11}^{q^2} - \nu_{11} + q_1(x) \cdot \nu_{21} = 0$, $\nu_{12}^{q^2} - \nu_{12} + q_1(x) \cdot \nu_{22} = 0$. We can set $\nu_{21} = \nu_{22} = 1$ and $\nu_{11} - \nu_{12} = 1$.

In this case,

$$(\tilde{\beta})^{-1}(Z_2^{\infty+0}) = \Delta$$

(Δ denotes the diagonal correspondence) because there exists a morphism of level structures between two vector bundles with the same degree and with ∞D -level structures. This morphism must be an isomorphism (Proposition 2.2).

As before, by Theorem 3.10 and by the explicit calculations in Remark 9 and Lemma 3.9 $\tilde{\beta}$ has associated the 2×2 -matrix:

$$\begin{pmatrix} 1 \otimes \alpha_{1,0}^0 & 1 \otimes (\alpha_{1,0}^0)^q \\ 1 \otimes \alpha_{2,0}^0 & 1 \otimes (\alpha_{2,0}^0)^q \end{pmatrix}^{-1} \cdot \begin{pmatrix} \alpha_{1,0}^0 \otimes 1 & (\alpha_{1,0}^0)^q \otimes 1 \\ \alpha_{2,0}^0 \otimes 1 & (\alpha_{2,0}^0)^q \otimes 1 \end{pmatrix} \times \\ \begin{pmatrix} 1 \otimes \nu_{11} & 1 \otimes \nu_{12} \\ 1 \otimes 1 & 1 \otimes 1 \end{pmatrix}^{-1} \cdot \begin{pmatrix} \nu_{11} \otimes 1 & \nu_{12} \otimes 1 \\ 1 \otimes 1 & 1 \otimes 1 \end{pmatrix}$$

If we denote by

$$A_{0,1}^0 := \begin{pmatrix} \alpha_{1,0}^0 \otimes 1 & (\alpha_{1,0}^0)^q \otimes 1 \\ \alpha_{2,0}^0 \otimes 1 & (\alpha_{2,0}^0)^q \otimes 1 \end{pmatrix}, A_{0,1}^1 := \begin{pmatrix} 1 \otimes \alpha_{1,0}^0 & 1 \otimes (\alpha_{1,0}^0)^q \\ 1 \otimes \alpha_{2,0}^0 & 1 \otimes (\alpha_{2,0}^0)^q \end{pmatrix}$$

and

$$A_\infty^0 := \begin{pmatrix} \nu_{11} \otimes 1 & \nu_{12} \otimes 1 \\ 1 \otimes 1 & 1 \otimes 1 \end{pmatrix}, A_\infty^1 := \begin{pmatrix} 1 \otimes \nu_{11} & 1 \otimes \nu_{12} \\ 1 \otimes 1 & 1 \otimes 1 \end{pmatrix}$$

then by Theorem 3.10:

$$(\tilde{\beta})^{-1}(Z_2^{\infty+0}) = (A_\infty^1 (A_{0,1}^1)^{-1} \cdot A_{0,1}^0 \cdot (A_\infty^0)^{-1} - Id_2 = 0)$$

Example. 4 Now, $n = 2$, $p(t) = t(t - 1)$. Thus, $D = 0 + 1$ and as in the above example we follow the notation of Remark 9:

$$\tilde{K}_2^{\infty+0} = \mathbb{F}_q(x)[\alpha_{1,0}^0, \alpha_{2,0}^0, \beta_{1,0}^0, \beta_{2,0}^0, \nu_{11}, \nu_{21}, \nu_{12}, \nu_{22}],$$

where $\alpha_{i,0}^0$ is as in the above example and

$$(\beta_{i,0}^0)^{q^2} + q_1(x) \cdot (\beta_{i,0}^0)^q + \beta_{i,0}^0 \cdot (x - 1) = 0$$

for $i = 1, 2$ and $(\nu_{ij})_{ij}$ as before.

A_∞^0 , A_∞^1 , $A_{0,1}^0$ and $A_{0,1}^1$ follow the same notation as in the last example and $B_{0,1}^0$, and $B_{0,1}^1$ denote

$$\begin{pmatrix} \beta_{1,0}^0 \otimes 1 & (\beta_{1,0}^0)^q \otimes 1 \\ \beta_{2,0}^0 \otimes 1 & (\beta_{2,0}^0)^q \otimes 1 \end{pmatrix}, \begin{pmatrix} 1 \otimes \beta_{1,0}^0 & 1 \otimes (\beta_{1,0}^0)^q \\ 1 \otimes \beta_{2,0}^0 & 1 \otimes (\beta_{2,0}^0)^q \end{pmatrix}$$

respectively.

Again by Theorem 3.10 and by the explicit calculations in Remark 9 and Lemma 3.9, $\tilde{\beta}$ has associated the 2×2 -matrix:

$$(A_\infty^1)^{-1} \cdot A_{0,1}^0 \times (B_\infty^1)^{-1} \cdot B_{0,1}^0 \times (A_\infty^1)^{-1} \cdot A_\infty^0$$

and $(\tilde{\beta})^{-1}(Z_2^{\infty+0+1})$ is

$$(A_\infty^1 (A_{0,1}^1)^{-1} \cdot A_{0,1}^0 \cdot (A_\infty^0)^{-1} - A_\infty^1 (B_{0,1}^1)^{-1} \cdot B_{0,1}^0 \cdot (A_\infty^0)^{-1} - Id_2 = 0).$$

The coefficients $+$ and $-$ are deduced bearing in mind the Chinese remainder theorem. We shall now attempt to calculate the irreducible components of $(\tilde{\beta})^{-1}(Z_2^{\infty+0+1})$. From Lemma 3.11, we know that these components are graphs $\Gamma_{g.F^i}$, where $i \leq 2$ and $g \in Gl_2(\mathbb{F}_q)_\infty \times Gl_2(\mathbb{F}_q)_0 \times Gl_2(\mathbb{F}_q)_1$.

We know from Lemma 3.12 that the graphs Γ_{F^2} and $\Gamma_{h_{d+t}}$ are contained in $(\tilde{\beta})^{-1}(Z_2^{\infty+0+1})$.

We try to prove that

$$(\tilde{\beta}_2^{\infty+0+1})^{-1}(\Theta_2^{\infty+0+1}) = \bigcup_{d \neq 0} \Gamma_{h_{d+t}} \cup \Gamma_{F^2}.$$

Let $g = g_\infty \times g_{0+1}$ be $B \times C \times D \in Gl_2(\mathbb{F}_q)_\infty \times Gl_2(\mathbb{F}_q)_0 \times Gl_2(\mathbb{F}_q)_1$. If $\Gamma_g \subset (\tilde{\beta})^{-1}(Z_2^{\infty+0+1})$, we have

$$(A_{0,1})^{-1} \cdot C \cdot A_{0,1} - (B_{0,1})^{-1} \cdot D \cdot B_{0,1} = (A_\infty)^{-1} \cdot B \cdot A_\infty$$

where

$$B_{0,1} := \begin{pmatrix} \beta_{1,0}^0 & (\beta_{1,0}^0)^q \\ \beta_{2,0}^0 & (\beta_{2,0}^0)^q \end{pmatrix} A_{0,1} := \begin{pmatrix} \alpha_{1,0}^0 & (\alpha_{1,0}^0)^q \\ \alpha_{2,0}^0 & (\alpha_{2,0}^0)^q \end{pmatrix}$$

and

$$A_\infty := \begin{pmatrix} \nu_{11} & \nu_{12} \\ 1 & 1 \end{pmatrix}.$$

By acting on this equality by $Id \times \Omega \times Id \in Gl_2(\mathbb{F}_q)_\infty \times Gl_2(\mathbb{F}_q)_0 \times Gl_2(\mathbb{F}_q)_1$, we obtain the expression

$$\begin{aligned} (A_{0,1})^{-1} \cdot \Omega^{-1} \cdot C \cdot \Omega \cdot A_{0,1} - (B_{0,1})^{-1} \cdot D \cdot B_{0,1} &= \\ &= (A_\infty)^{-1} \cdot B \cdot A_\infty. \end{aligned}$$

Thus, for each $\Omega \in Gl_2(\mathbb{F}_q)$ we have that $\Omega^{-1} \cdot C \cdot \Omega = C$, and hence C is a central matrix. In an analogous way, we can check that B and D must be central matrices. As $\mathbb{F}_q^*$ acts on $\tilde{K}_2^{\infty+0+1}$ by the identity, we can assume that $B = Id$. Then, $C - D = Id$, and hence $g = h_{d+t}$.

In a similar way, one can check that there does not exist g with

$$\Gamma_{g.F} \subset (\tilde{\beta})^{-1}(Z_2^{\infty+0+1}).$$

We thus obtain what were looking for:

$$(\tilde{\beta})^{-1}(Z_2^{\infty+0+1}) = \bigcup_{d \neq 0} \Gamma_{h_{d+t}} \cup \Gamma_{F^2}.$$

4. CASE OF A GENERAL CURVE

In this section we shall study zeta subschemes for a general curve C . We shall translate the results of the above sections to this case. Now, instead of ∞ we shall consider a rational point $p \in \text{Spec}(A)$ such that $p \notin \text{Supp}(D)$. In this case, we shall state in a more general way the result of Lemma 3.11, and the relation between these zeta subschemes and the general theta divisor.

Let us recall some notation and results: $\mathcal{M}_C(n, h)$ denotes the moduli stack, [LM], of vector bundles of rank n and degree h . In the same way, $\mathcal{M}_C(n, h, p + D)$ denotes the moduli stack of pairs, (M, f_{p+D}) , of vector bundles of rank n and degree h with a $p + D$ -level structure. There exists a natural epimorphism of forgetting the level structures:

$$L : \mathcal{M}_C(n, h, p + D) \rightarrow \mathcal{M}_C(n, h)$$

Let us denote by $\Theta_n \subset \mathcal{M}_C(n, n(g-1))$ the general theta divisor [DN]: Θ_n is the closed substack of $\mathcal{M}_C(n, n(g-1))$ of vector bundles M such that $H^0(C, M) = 0$. We set $\Theta_n^{\infty D} := L^{-1}(\Theta_n)$.

By $\mathcal{E}_n^{p+D}$ we denote the moduli (scheme for $\deg(D) \gg 0$) of rank n - A -elliptic sheaves, (E_j, i_j, τ) , with $p + D$ -level structures. As in the preceding sections, we have a natural morphism

$$\Phi : \mathcal{E}_n^{p+D} \rightarrow \mathcal{M}_C(n, ng, p + D),$$

given by $\Phi(E_j, i_j, \tau, f_{p+D}) = (E_0, f_{p+D}^0)$.

In this section, π denotes the natural morphism $\mathcal{O}_C(g+d-1) \rightarrow \mathcal{O}_C/\mathcal{O}_C(-p-D)$. As in the above sections

$$\mathcal{O}_{pD} := \mathcal{O}_C/\mathcal{O}_C(-p-D) \text{ and } \mathcal{O}_{pD} := H^0(C, \mathcal{O}_C/\mathcal{O}_C(-p-D)).$$

Remark 10. We consider the open substack

$$\mathcal{U}_d \subseteq \mathcal{M}_C(n, h, p + D) \times \mathcal{M}_C(n, h, p + D)$$

of pairs $[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})]$ satisfying $H^1(C, \bar{M}^\vee \otimes_{\mathcal{O}_C} M(g+d-1)) = 0$. In this case, if $[(\bar{\mathbb{M}}, \bar{f}_{p+D}), (\mathbb{M}, f_{p+D})]$ is a universal object for $\mathcal{U}_d$ we have

$$h^0(\bar{\mathbb{M}}^\vee \otimes_{\mathcal{O}_C \times \mathcal{U}_d} \mathbb{M}(g+d-1)) = n^2 d.$$

Let $\{s_1, \dots, s_{n^2 d}\}$ be a basis for the space global sections of

$$\bar{\mathbb{M}}^\vee \otimes_{\mathcal{O}_C \times \mathcal{U}_d} \mathbb{M}(g+d-1),$$

and we set

$$o_i = H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi)(s_i).$$

In this last Remark and in the following Lemmas we assume, for easy notation, that the global sections of $\bar{\mathbb{M}}^\vee \otimes_{\mathcal{O}_C \times \mathcal{U}_d} \mathbb{M}(g+d-1)$ are a free module of rank $n^2 d$. By

Grauert's Theorem, it occurs locally, so instead of $\mathcal{U}_d$ we should write $\mathcal{U}_d^\alpha$, where $\mathcal{U}_d = \bigcup_\alpha \mathcal{U}_d^\alpha$. The localization by $\mathcal{U}_d^\alpha$ of the space of global sections of

$$\bar{M}^\vee \otimes_{\mathcal{O}_C \otimes \mathcal{O}_{\mathcal{U}_d}} M(g+d-1)$$

is a $\mathcal{O}_{\mathcal{U}_d^\alpha}$ -free module.

We shall now define an analogous subscheme to $Z_n^{\infty D}$.

Definition 4.1. *Let S be a scheme. We define the $p+D$ -generalized zeta substack, Z_n^{p+D} , as the closed substack of $\mathcal{U}_d$ defined by: $[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in \text{Hom}_{\text{stacks}}(S, Z_n^{p+D})$ if over the open subscheme, $\bar{U}$, of S formed by the $s \in S$ such that*

$$H^0(C \otimes k(s), \bar{M}_s^\vee \otimes_{\mathcal{O}_{C \times k(s)}} M_s((g+d-1)\infty - p - D)) = 0$$

there exists a morphism of modules $T : \bar{M}|_{\bar{U}} \rightarrow M(g+d-1)|_{\bar{U}}$ ($d = \deg(D)$), where the diagram :

$$\begin{array}{ccc} \bar{M}|_{\bar{U}} & \xrightarrow{T} & M(g+d-1)|_{\bar{U}} \\ & \searrow \bar{f}_{p+D} & \downarrow f_{p+D} \otimes \pi \\ & & (\mathcal{O}_{pD})^n \otimes \mathcal{O}_{\bar{U}} \end{array}$$

is commutative.

Remark 11. From the last Lemma, if

$$[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in \text{Hom}_{\text{stacks}}(S, Z_n^{p+D})$$

then there exists an open subscheme $\bar{U} \subset S$ such that over this open subscheme there exists a non-trivial morphism $T : \bar{M} \rightarrow M(g+d-1)$. And if $s \notin \bar{U}$, there exists a non-trivial morphism $T_s : \bar{M}_s \rightarrow M_s((g+d-1)\infty - p - D) \subset M(g+d-1)_s$.

Hence $Z_n^{p+D} = V_n^{p+D} \cup (V_n^{p+D})^c$, with V_n^{p+D} an open substack of Z_n^{p+D} .

Remark 12. A pair $[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in \text{Hom}_{\text{stacks}}(\text{Spec}(R), Z_n^{p+D})$ gives a morphism

$$\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi : \bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} M(g+d-1) \longrightarrow (\mathcal{O}_{pD})^{n^2} \otimes R.$$

Therefore, by taking global sections we obtain a morphism

$$H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi) : H^0(\mathcal{O}_C \otimes R, \bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} M(g+d-1)) \rightarrow (\mathcal{O}_{pD})^{n^2} \otimes R.$$

We shall denote by Ξ the natural morphism

$$\Xi : \mathcal{O}_C \otimes R \rightarrow \bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} \bar{M} \xrightarrow{\bar{f}_{p+D}^\vee \otimes \bar{f}_{p+D}} (\mathcal{O}_{pD})^{n^2} \otimes R$$

where the first morphism in the composition is the diagonal morphism. Therefore, by taking global sections we obtain a morphism

$$H^0(\Xi) : R \rightarrow (\mathcal{O}_{pD})^{n^2} \otimes R$$

given by the diagonal matrix with entries over $\mathcal{O}_{pD}$.

Let us now consider the morphism

$$\Sigma := H^0(\Xi) + H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi)$$

from $R \oplus H^0(\mathcal{O}_C \otimes R, \bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} M(g+d-1))$ to $(\mathcal{O}_{pD})^{n^2} \otimes R$.

Lemma 4.2. A pair $[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in \text{Hom}_{\text{stacks}}(\text{Spec}(R), \mathcal{U}_d)$ belongs to $\text{Hom}_{\text{stacks}}(\text{Spec}(R), Z_n^{p+D})$

if and only if Σ is not injective.

Proof. Since if $s \in \text{Spec}(R)$ satisfies

$$H^0(C \otimes k(s), \bar{M}_s^\vee \otimes_{\mathcal{O}_{C \times k(s)}} M_s((g+d-1)\infty - p - D)) \neq 0$$

we have $\text{Ker}(H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi)_s) \neq 0$. Thus, Σ_s is not injective and we can assume that for all $s \in \text{Spec}(R)$ we have:

$$H^0(C \otimes k(s), \bar{M}_s^\vee \otimes_{\mathcal{O}_{C \times k(s)}} M_s((g+d-1)\infty - p - D)) = 0.$$

If there exists a morphism T between the level structures considered

$$\begin{array}{ccc} \bar{M} & \xrightarrow{T} & M(g+d-1) \\ & \searrow \bar{f}_{p+D} & \downarrow f_{p+D} \otimes \pi \\ & & (\mathcal{O}_{pD})^n \otimes R \end{array}$$

by tensoring by $\bar{M}^\vee$ and considering $\bar{f}_{p+D}$, we obtain a section

$$w \in H^0(\mathcal{O}_C \otimes R, \bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} M(g+d-1))$$

such that

$$H^0(\Xi)(1) = H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi)(w).$$

Thus, in this case $1 \oplus -w \in \text{Ker}(\Sigma)$.

Conversely, if $1 \oplus -s \in \text{Ker}(\Sigma)$ we have a commutative diagram

$$\begin{array}{ccc} \mathcal{O}_C \otimes R & \longrightarrow & \bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} M(g+d-1) \\ & \searrow \Xi & \downarrow \bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi \\ & & (\mathcal{O}_{pD})^{n^2} \otimes R \end{array}$$

Therefore, we conclude by tensoring by $\bar{M}$ and bearing in mind $\bar{f}_{p+D}$. □

Remark 13. If $Z_n^{p+D} = V_n^{p+D} \cup (V_n^{p+D})^c$, we deduce that that:

$$(V_n^{p+D})^c = \{[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in Z_n^{p+D}, \text{ with } \text{Ker}(H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi)) \neq 0\}$$

and $[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in V_n^{p+D}$ if

$$H^0(\Xi)(1) \in H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi).$$

Remark 14. From $H^0(\Xi)(1)$ and $\{o_1, \dots, o_{n^2d}\}$, we obtain a $(n^2d+1)(\text{columns}) \times (n^2(d+1))(\text{rows})$ -matrix $(H^0(\Xi)(1), o_1, \dots, o_{n^2d})$ with entries over $H^0(\mathcal{U}_d, \mathcal{O}_{\mathcal{U}_d})$. Recall that $\{H^0(\Xi)(1), o_1, \dots, o_{n^2d}\} \subset (\mathcal{O}_{pD})^n \otimes H^0(\mathcal{U}_d, \mathcal{O}_{\mathcal{U}_d})$ and $\dim_{\mathbb{F}_q} \mathcal{O}_{pD} = d+1$.

Let $\bar{\mathcal{U}}_d$ be the open substack of $\mathcal{U}_d$ such that $[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in \bar{\mathcal{U}}_d$ if and only if $h^0(\bar{M}^\vee \otimes_{\mathcal{O}_C} M(g+d-1)(-p-D)) = 0$. In this case, $\text{rank}(o_1, \dots, o_{n^2d}) = n^2d$.

Lemma 4.3. Z_n^{p+D} is a closed substack of $\mathcal{U}_d$ and is formed by the pairs

$$[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in \mathcal{U}_d,$$

such that

$$\text{rank}((O_{p+D})^{n^2} / < H^0(\Xi)(1), o_1, \dots, o_{n^2d} >) \geq n^2.$$

Moreover, $V_n^{p+D} = \bar{\mathcal{U}}_d \cap Z_n^{p+D}$ is locally the zero locus of n^2 -regular functions.

Proof. The first assertion is easily deduced from Lemma 4.2.

For the second assertion, we denote by

$$N_{i_1, \dots, i_{n^2d+1}}^{(H^0(\Xi)(1), o_1, \dots, o_{n^2d})}$$

$(1 \leq i_1 < \dots < i_{n^2d+1} \leq n^2(d+1))$ the n^2d+1 -minor of the matrix

$$(H^0(\Xi)(1), o_1, \dots, o_{n^2d}),$$

where the k^{th} -row is the i_k^{th} -row of $(H^0(\Xi)(1), o_1, \dots, o_{n^2d})$.

Analogously, we set

$$N_{i_1, \dots, i_{n^2d}}^{(o_1, \dots, o_{n^2d})}$$

for the $(n^2d)(\text{columns}) \times (n^2(d+1))(\text{rows})$ -matrix $(o_1, \dots, o_{n^2d})$. Let us take the open covering of $\bar{\mathcal{U}}_d$

$$\bar{\mathcal{U}}_d = \bigcup_{1 \leq i_1 < \dots < i_{n^2d} \leq n^2(d+1)} \bar{\mathcal{U}}_d^{i_1, \dots, i_{n^2d}}$$

$\bar{\mathcal{U}}_d^{i_1, \dots, i_{n^2d}}$ being the open substack of $\bar{\mathcal{U}}_d$ of pairs $[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in \mathcal{U}_d$, such that

$$\text{rank}(N_{i_1, \dots, i_{n^2d}}^{(o_1, \dots, o_{n^2d})}) = \text{rank}(o_1, \dots, o_{n^2d}) = n^2d.$$

Thus, we have that $\bar{\mathcal{U}}_d^{i_1, \dots, i_{n^2d}} \cap Z_n^{p+D}$ is the zero locus of the functions

$$\{\det(N_{i_1, \dots, i_{n^2d}, j_{n^2d+1}}^{(H^0(\Xi)(1), o_1, \dots, o_{n^2d})}), \dots, \det(N_{i_1, \dots, i_{n^2d}, j_{n^2d+n^2}}^{(H^0(\Xi)(1), o_1, \dots, o_{n^2d})})\},$$

with $\{1, 2, \dots, n^2(d+1)\} \setminus \{i_1, \dots, i_{n^2d}\} = \{j_{n^2d+1}, \dots, j_{n^2d+n^2}\}$.

□

If

$$[(\bar{M}, \bar{f}_{p+D}), (M, f_{p+D})] \in \text{Hom}_{\text{stacks}}(\text{Spec}(R), \bar{\mathcal{U}}_d),$$

we have

$$H^0(C \otimes R, \bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} M(g+d-1)(-p-D)) = 0$$

and

$$H^1(C \otimes R, \bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} M(g+d-1)) = 0.$$

In this way,

$$H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi)(H^0(C \otimes R, \bar{M}^\vee \otimes_{\mathcal{O}_C} M(g+d-1)))$$

is a rank n^2d -sub-vector bundle of $(O_{pD})^{n^2} \otimes R$. Thus,

$$\bigwedge^{n^2d} (H^0(\bar{f}_{p+D}^\vee \otimes f_{p+D} \otimes \pi)(H^0(\bar{M}^\vee \otimes_{\mathcal{O}_C \otimes R} M(g+d-1)))) \in \mathbb{P}(\bigwedge^{n^2d} (O_{pD})^{n^2} \otimes R)$$

and we obtain a morphism

$$\bar{\mathcal{U}}_d \rightarrow \mathbb{P}(\bigwedge^{n^2 d} (O_{pD})^{n^2}).$$

It is not hard to prove that

$$\det(N_{i_1, \dots, i_{n^2 d}, j_{n^2 d+k}}^{(H^0(\Xi)(1), o_1, \dots, o_{n^2 d})})$$

can be considered as the restriction to $\bar{\mathcal{U}}_d$ of a certain hyperplane section of $\mathbb{P}(\bigwedge^{n^2 d} (O_{pD})^{n^2})$. In this way, $\mathcal{U}_d^{i_1, \dots, i_{n^2 d}} \cap Z_n^{p+D}$ is obtained as the intersection of n^2 hyperplane sections of $\mathbb{P}(\bigwedge^{n^2 d} (O_{pD})^{n^2})$.

For $n = 1$, we have a direct relation of Z_1^{p+D} with a theta divisor. We take $d > 2g - 2$.

Definition 4.4. We define the $p+D$ -generalized theta divisor, Θ_{p+D} , as the closed subscheme of $J_{C,0}^{p+D}$ defined by: $(L, f_{p+D}) \in \text{Hom}_{\text{schemes}}(\text{Spec}(K), \Theta_{p+D})$ (K a field) if and only if there exists a morphism of modules $T : \mathcal{O}_C \otimes R \rightarrow L(g+d-1)$ ($d = \deg(D)$), where the diagram:

$$\begin{array}{ccc} \mathcal{O}_C \otimes K & \xrightarrow{T} & L(g+d-1) \\ \downarrow & \swarrow f_{p+D} \otimes \pi & \\ \mathcal{O}_{p+D} \otimes K & & \end{array}$$

is commutative. The vertical arrow is the natural epimorphism.

Let us consider $J_{C,g}^{p+D}$, where $J_{C,g}^{p+D}$ is the $p+D$ -generalized Rosenlich's Jacobian (c.f:[Se]).

Remark 15. By regarding the morphism

$$m_{p+D} : J_{C,g}^{p+D} \times J_{C,g}^{\infty D} \rightarrow J_{C,0}^{p+D}$$

defined by

$$m_{p+D}[(\bar{L}, \bar{f}_{p+D}), (L, f_{p+D})] = (\bar{L}^\vee, \bar{f}_{p+D}^\vee) \otimes (L, f_{p+D}),$$

we have that $m_{\infty D}^{-1}(\Theta_1^{\infty D}) = Z_1^{\infty D}$ for $d > 2g - 2$, because of the definition of $Z_1^{\infty D}$ and $\mathcal{U}_d = J_{C,g}^{p+D} \times J_{C,g}^{\infty D}$.

Moreover, from the last Lemma we can say that $\bar{\mathcal{U}}_d^{i_1, \dots, i_d} \cap Z_1^{p+D}$ is the zero locus of the function

$$\det(N_{i_1, \dots, i_d, j_{d+1}}^{(1, o_1, \dots, o_d)})$$

over $\bar{\mathcal{U}}_d^{i_1, \dots, i_d}$ (for $n = 1$ $H^0(\Xi)(1) = 1 \in O_{pD}$). For different $\{i_1, \dots, i_d\}$, these determinants are the determinant $\det(1, o_1, \dots, o_d)$ up to a sign. Then,

$$\bar{\mathcal{U}}_d \cap Z_1^{p+D} = \bar{\mathcal{U}}_d \cap (\det(1, o_1, \dots, o_d))_0.$$

Recall that

$$\bar{\mathcal{U}}_d = \bigcup_{1 \leq i_1 < \dots < i_{n^2 d} \leq d+1} \bar{\mathcal{U}}_d^{i_1, \dots, i_{n^2 d}}.$$

On the other hand, as $(\bar{\mathcal{U}}_d)^c$ is formed by the pairs where $\{o_1, \dots, o_d\}$ are linearly dependent, then

$$(\bar{\mathcal{U}}_d)^c \subseteq (\det(1, o_1, \dots, o_d))_0,$$

and we conclude that $Z_1^{p+D} = (\det(1, o_1, \dots, o_d))_0$.

Recall that actually this result is settled locally over $J_{C,g}^{p+D} \times J_{C,g}^{p+D}$. To be precise, one must take a universal object $[(\bar{\mathbb{L}}, \bar{f}), (\mathbb{L}, f)]$ and an open covering for $J_{C,g}^{p+D} \times J_{C,g}^{p+D}$ such that

$$p_*(\bar{\mathbb{L}}^\vee \otimes_{\mathcal{O}_C \otimes \mathcal{O}_J} \mathbb{L})$$

is trivialized by this covering. p denotes the natural projection $C \times J_{C,g}^{p+D} \rightarrow J_{C,g}^{p+D}$.

4.1. Zeta subschemes for rank- n -Shtukas. In this section we shall study zeta subschemes in the case of rank- n -Shtukas and A -Drinfeld modules defined over an arbitrary curve C . Professor G.W. Anderson has suggested that I should study this topic this topic.

First, we shall state some previous definitions.

Definition 4.5. *A rank- n shtuka over $C \times \text{Spec}(R)$ is a diagram of rank- n -vector bundles over $C \times \text{Spec}(R)$*

$$\mathcal{L}_r \xrightarrow{j} \mathcal{L}_{r+1} \xleftarrow{t} F^\# \mathcal{L}_r,$$

such that $\mathcal{L}_{r+1}/j(\mathcal{L}_r)$ and $\mathcal{L}_{r+1}/t(F^\# \mathcal{L}_r)$ are line bundles as R -modules and they are supported by two morphisms Γ_α and Γ_γ $\text{Spec}(R) \rightarrow C$; the pole and zero, respectively. We denote the line bundles associated with Γ_α and Γ_γ by $\mathcal{O}_{C \otimes R}(\alpha)$ and $\mathcal{O}_{C \otimes R}(\gamma)$ respectively. We assume $j(\mathcal{L}_r) + t(F^\# \mathcal{L}_r) = \mathcal{L}_{r+1}$. We set $\deg(\mathcal{L}_r) = ng + r$.

Definition 4.6. *Similar to the above sections, to give a $p + D$ -level structure over*

$$\mathcal{L}_r \xrightarrow{j} \mathcal{L}_{r+1} \xleftarrow{t} F^\# \mathcal{L}_r$$

is to give level structures $(\mathcal{L}_r, f_{p+D}^r)$ and $(\mathcal{L}_{r+1}, f_{p+D}^{r+1})$ with commutative diagrams

$$\begin{array}{ccc} \mathcal{L}_r & \xrightarrow{j} & \mathcal{L}_{r+1} \\ & \searrow f_{p+D}^r & \downarrow f_{p+D}^{r+1} \\ & & (\mathcal{O}_{p+D})^n \otimes R \end{array}$$

and

$$\begin{array}{ccc} F^\# \mathcal{L}_r & \xrightarrow{t} & \mathcal{L}_{r+1} \\ & \searrow F^\# f_{p+D}^r & \downarrow f_{p+D}^{r+1} \\ & & (\mathcal{O}_{p+D})^n \otimes R \end{array},$$

$p \notin \{\alpha, \gamma, \text{supp}(D)\}$.

Definition 4.7. *An isogeny between two rank- n Shtukas*

$$\mathcal{L}_r \xrightarrow{j} \mathcal{L}_{r+1} \xleftarrow{t} F^\# \mathcal{L}_r$$

and

$$\mathcal{N}_h \xrightarrow{j'} \mathcal{N}_{h+1} \xleftarrow{t'} F^\# \mathcal{N}_h$$

is defined by two morphisms of $\mathcal{O}_{C \otimes R}$ -modules $T : \mathcal{L}_r \rightarrow \mathcal{N}_h$ and $H : \mathcal{L}_{r+1} \rightarrow \mathcal{N}_{h+1}$ such that $j'.T = H.j$ and $t'.F^\#T = F^\#H.t$.

Analogously, one can define a $p + D$ -isogeny between shtukas with $p + D$ -level structures, except that one must take T and H compatible with the level structures.

We now consider the stack, $\text{Cht}_{p+D}^{n, ng+r}$, of shtukas of rank n and $\deg(\mathcal{L}_r) = ng+r$. We follow the same notation for $\text{Cht}_{p+D}^{n, ng+r}$ but considering $p + D$ -level structures. Let $P^{n, ng+r}$ be the closed substack of $\text{Cht}_{p+D}^{n, ng+r} \times \text{Cht}_{p+D}^{n, ng+r}$ of shtukas with the same pole. Analogously, we define $P_{p+D}^{n, ng+r}$.

Proposition 4.8. Let $[(\mathcal{L}_r, \mathcal{L}_{r+1}, j, t), (\mathcal{N}_h, \mathcal{N}_{h+1}, j', t')]$ be an element of $P^{n, ng+r}$ with pole α and defined over a $\mathbb{F}_q$ -reduced algebra R . Then, there exists an isogeny between these shtukas if and only if there exists a morphism of modules $T : \mathcal{L}_r \rightarrow \mathcal{N}_h$, together with the commutative diagram

$$\begin{array}{ccc} \mathcal{L}_r(\alpha) & \xrightarrow{T(\alpha)} & \mathcal{N}_h(\alpha) \\ \uparrow \kappa & & \uparrow \kappa' \\ \mathcal{L}_{r+1} & & \mathcal{N}_{h+1} \\ \uparrow t & & \uparrow t' \\ F^\# \mathcal{L}_r & \xrightarrow{F^\# T} & F^\# \mathcal{N}_h \end{array}$$

$T(\alpha)$ denotes the morphism induced by T from $\mathcal{L}_r(\alpha)$ to $\mathcal{N}_h(\alpha)$. The injective morphisms $\kappa : \mathcal{L}_{r+1} \hookrightarrow \mathcal{L}_r(\alpha)$ and $\kappa' : \mathcal{N}_{h+1} \hookrightarrow \mathcal{N}_h(\alpha)$ are morphisms such that the composition with j and j' are the morphisms $\mathcal{L}_r \hookrightarrow \mathcal{L}_r(\alpha)$ and $\mathcal{N}_h \hookrightarrow \mathcal{N}_h(\alpha)$ induced by the natural inclusion $\mathcal{O}_C \otimes R \subset \mathcal{O}_C \otimes R(\alpha)$. Recall that the cokernels of $\mathcal{L}_r \hookrightarrow \mathcal{L}_{r+1}$ and $\mathcal{N}_h \hookrightarrow \mathcal{N}_{h+1}$ are supported over α .

Proof. This can be deduced because

$$\begin{array}{ccc} \mathcal{L}_r(\alpha) & \xrightarrow{T(\alpha)} & \mathcal{N}_h(\alpha) \\ \uparrow \kappa & & \uparrow \kappa' \\ \mathcal{L}_{r+1} & & \mathcal{N}_{h+1} \\ \uparrow j & & \uparrow j' \\ \mathcal{L}_r & \xrightarrow{T} & \mathcal{N}_h \end{array}$$

is commutative, and

$$j(\mathcal{L}_r) + t(F^\# \mathcal{L}_r) = \mathcal{L}_{r+1}, \quad j'(\mathcal{N}_h) + t'(F^\# \mathcal{N}_h) = \mathcal{N}_{h+1}.$$

□

Let consider us the natural morphism $\bar{\Phi} : \text{Cht}_{p+D}^{n, ng+r} \rightarrow \mathcal{M}_C(n, ng+r, p+D)$ given by $\bar{\Phi}((\mathcal{L}_r, f_{p+D}^r), (\mathcal{L}_{r+1}, f_{p+D}^{r+1}), j, t) = (\mathcal{L}_r, f_{p+D}^r)$. Hence, we deduce a morphism

$$\bar{\beta} : P_{p+D}^{n, ng+r} \rightarrow \text{Cht}_{p+D}^{n, ng+r} \times \text{Cht}_{p+D}^{n, ng+r} \xrightarrow{\bar{\Phi} \times \bar{\Phi}} \mathcal{M}_C(n, ng+r, p+D) \times \mathcal{M}_C(n, ng+r, p+D).$$

For easy notation, we denote by $(\mathcal{L}_r, \mathcal{L}_{r+1}, f_{p+D})$ an element of $\text{Cht}_{p+D}^{n, ng+r}$. Because $p \neq \alpha$ in the following Lemma, we are able to set, for shtukas, the results of

Lemma 3.7 in a more precise way. This Lemma attempts to be an approximation for shtukas with level structures of Lemma 3.3 [An2].

Lemma 4.9. Let $[(\mathcal{L}_0, \mathcal{L}_1, f_{p+D}), (\mathcal{N}_0, \mathcal{N}_1, g_{p+D})] \in \bar{\beta}^{-1}(Z_n^{p+D})$ be defined over a field K . Then:

1) There exists a $p+D$ -isogeny between the shtukas with $p+D$ -level structures $[(\mathcal{L}_0, \mathcal{L}_1, f_{p+D})$ and $(\mathcal{N}_0, \mathcal{N}_1, g_{p+D} \otimes \pi)]$.

or

2) There exists an isogeny between the shtuka $(\mathcal{L}_0, \mathcal{L}_1, j, t)$ and the twisted shtuka $(\mathcal{N}_0, \mathcal{N}_1, j', t') \otimes_{\mathcal{O}_C} ((g+d-1)\infty - p - D)$.

or

3) $(F^\# \mathcal{L}_0)^\vee \otimes_{\mathcal{O}_C \otimes K} \mathcal{N}_0((g+d-1)\infty - p - D + \alpha) \in \Theta_{n^2}$.

Proof. Let us consider the diagram (not necessarily commutative)

$$\begin{array}{ccccc}
 \mathcal{L}_0(\alpha) & \xrightarrow{T(\alpha)} & \mathcal{N}_0(g+d-1)(\alpha) & & \\
 \uparrow \kappa & \searrow & \swarrow & \uparrow \kappa' & \\
 \mathcal{L}_1 & & (\mathcal{O}_{p+D})^n \otimes R & & \mathcal{N}_1(g+d-1) \\
 \uparrow t & \swarrow & \nwarrow & \uparrow t' & \\
 F^\# \mathcal{L}_0 & \xrightarrow{F^\# T} & F^\# \mathcal{N}_0(g+d-1) & &
 \end{array}$$

T is a morphism given by $[(\mathcal{L}_0, f_{p+D}^0), (\mathcal{N}_0, g_{p+D}^0)] \in Z_n^{p+D}$. The diagonal morphisms are the level structures and the vertical morphisms κ and κ' are induced by j and j' since these two shtukas have the same pole, α . From Definition 4.1, we have two possibilities: either $T : \mathcal{L}_0 \rightarrow \mathcal{N}_0(g+d-1)$ takes values in $\mathcal{N}_0(g+d-1)(-p-D)$ or the latter diagram is commutative after tensoring by $\mathcal{O}_{pD}$. In both cases, one obtains a morphism:

$$T(\alpha) \cdot \kappa \cdot t - \kappa' \cdot t' \cdot F^\# T : F^\# \mathcal{L}_0 \rightarrow \mathcal{N}_0(g+d-1)(-p-D+\alpha).$$

When $T(\alpha) \cdot \kappa \cdot t - \kappa' \cdot t' \cdot F^\# T = 0$, we obtain cases 1) and 2) and when it is $\neq 0$ it tell us that $(F^\# \mathcal{L}_0)^\vee \otimes_{\mathcal{O}_C \otimes R} \mathcal{N}_0((g+d-1)\infty - p - D + \alpha) \in \Theta_{n^2}$. $\square$

4.2. Zeta subschemes and A -Drinfeld modules. In this subsection, by means of the natural immersion of elliptic sheaves into shtukas (c.f.[Dr3]), we shall be able to translate the results of the above section to the case of Drinfeld modules. We use the notation and definitions of section 3.

Proposition 4.10. There exists an immersion of stacks

$$\Upsilon : \mathcal{E}_n^{p+D} \hookrightarrow \text{Ch}t_{p+D}^{n,ng}$$

Proof. This is defined by

$$\Upsilon(E_j, i_j, \tau, f_{p+D}) = ((E_0, f_{p+D}^0), (E_1, f_{p+D}^1), i_0, \tau)$$

This immersion is given in [Dr3] (1.3). $\square$

We recall the definition of isogeny for A -Drinfeld modules

Definition 4.11. An isogeny of degree r between two Drinfeld modules ϕ and $\bar{\phi}$ is a polynomial of degree r in σ , $q(\sigma)$, such that the endomorphism $q(\sigma) : (\mathbb{G}_a)_R \rightarrow (\mathbb{G}_a)_R$ satisfies

$$q(\sigma)(\phi_a(\lambda)) = \bar{\phi}_a(q(\sigma))(\lambda)$$

for all $a \in A$.

Definition 4.12. Two elliptic sheaves $(\bar{E}_j, \bar{i}_j, \bar{\tau})$ and (E_j, i_j, τ) are said to be isogenous by an isogeny of degree $\leq r$ if and only if there exist morphisms of modules $T_j : \bar{E}_j \rightarrow E_{j+r}$, for each j , compatible with the diagrams that define the elliptic sheaves.

Analogously, one can define a $p+D$ -isogeny between elliptic sheaves with $p+D$ -level structures, except that one must take " T_j " compatible with the level structures.

There exists an equivalence between the notion of isogenous elliptic sheaves and isogenous Drinfeld modules:

Proposition 4.13. Two Drinfeld modules ϕ and $\bar{\phi}$ associated with elliptic sheaves $(\bar{E}_j, \bar{i}_j, \bar{\tau})$ and (E_j, i_j, τ) , respectively, are isogenous by an isogeny of degree $\leq r$ if and only if there exists an isogeny of degree $\leq r$ between the elliptic sheaves $(\bar{E}_j, \bar{i}_j, \bar{\tau})$ and (E_{j+r}, i_{j+r}, τ) .

The following Proposition together with Proposition 4.8 tell us that two elliptic sheaves are isogenous as shtukas if and only if they are isogenous as elliptic sheaves. Recall that the elliptic sheaves considered have ∞ as their pole.

Proposition 4.14. Two elliptic sheaves $(\bar{E}_j, \bar{i}_j, \bar{\tau})$ and (E_j, i_j, τ) defined over a $\mathbb{F}_q$ -reduced algebra, R , are isogenous by an isogeny of degree $\leq r$ if and only if there exists a morphism of vector bundles

$$T : \bar{E}_0 \rightarrow E_r$$

with the diagram

$$\begin{array}{ccc} \bar{E}_0 & \xrightarrow{T} & E_r \\ \bar{\tau} \uparrow & & \uparrow \tau \\ F^\# \bar{E}_{-n} & \xrightarrow{F^\# T(-1)} & F^\# E_{r-n} \end{array}$$

commutative. Here, $T(-1)$ denotes the morphism induced by T over

$$\bar{E}_{-n} \rightarrow E_{r-n},$$

bearing in mind that $E_{k-n} := E_k(-\infty)$.

Proof. Since $\bar{E}_{1-n} = \bar{\tau}(F^\# \bar{E}_{-n}) + \bar{E}_{-n} \subset \bar{E}_0$ and $E_{r+1-n} = \tau(F^\# E_{r-n}) + E_{r-n} \subset E_r$, by observing the latter diagram we have $T(\bar{E}_{1-n}) \subset E_{r+1-n}$. Hence, we obtain the commutative diagram

$$\begin{array}{ccc} \bar{E}_0 & \xrightarrow{T} & E_r \\ \uparrow \bar{\tau} & & \uparrow \tau \\ \bar{E}_{1-n} & \xrightarrow{T_{1-n}} & E_{r+1-n} \\ \uparrow \bar{\tau} & & \uparrow \tau \\ \bar{E}_{-n} & \xrightarrow{T(-1)} & E_{r-n} \end{array} \quad ,$$

where T_{1-n} denotes the restriction of T to $\bar{E}_{1-n}$ and the injective morphisms are the " i " and " $\bar{i}$ " inclusions given in the elliptic sheaves.

From the last commutative diagram we also obtain a commutative diagram

$$\begin{array}{ccc} F^\# \bar{E}_{1-n} & \xrightarrow{F^\# T_{1-n}} & F^\# E_{r+1-n} \\ \uparrow \cup & & \uparrow \cup \\ F^\# \bar{E}_{-n} & \xrightarrow{F^\# T(-1)} & F^\# E_{r-n} \end{array}$$

and in the case of R being a field,

$$\begin{array}{ccc} \bar{E}_0 & \xrightarrow{T} & E_r \\ \bar{\tau} \uparrow & & \uparrow \tau \\ F^\# \bar{E}_{1-n} & \xrightarrow{F^\# T_{1-n}} & F^\# E_{r+1-n} \end{array} ,$$

($\bar{\tau}$ and τ are morphisms induced by the definition of elliptic sheaves) is also commutative because $(T.\bar{\tau} - \tau.F^\# T_{1-n})|_{F^\# \bar{E}_{-n}} = 0$. Then,

$$F^\# \bar{E}_{-n} \subset \text{Ker}(T.\bar{\tau} - \tau.F^\# T_{1-n})$$

and hence $\text{rank}(\text{Ker}(T.\bar{\tau} - \tau.F^\# T_{1-n})) = n$. Because $T.\bar{\tau} - \tau.F^\# T_{1-n}$ defines an injective morphism

$$F^\# \bar{E}_{1-n} / \text{Ker}(T.\bar{\tau} - \tau.F^\# T_{1-n}) \hookrightarrow E_{r-n},$$

we conclude that $T.\bar{\tau} - \tau.F^\# T_{1-n} = 0$. We finish the proof because for each $s \in \text{Spec}(R)$ we have $\text{Im}(T.\bar{\tau} - \tau.F^\# T_{1-n}) \subset m_s.E_r$ for all $s \in \text{Spec}(R)$ then, as E_r is locally free and R is reduced, we have $T.\bar{\tau} - \tau.F^\# T_{1-n} = 0$.

We repeat the same argument to obtain $T_{l-n} : \bar{E}_{l-n} \rightarrow E_{r+l-n}$ with $0 \leq l \leq n$. $\square$

By looking at Υ and at the morphism defined in the last subsection,

$$\bar{\beta} : P_{p+D}^{n,ng+r} \rightarrow \text{Ch}t_{p+D}^{n,ng+r} \times \text{Ch}t_{p+D}^{n,ng+r} \xrightarrow{\bar{\Phi} \times \bar{\Phi}} \mathcal{M}_C(n, ng+r, p+D) \times \mathcal{M}_C(n, ng+r, p+D),$$

we obtain, in the setting of elliptic sheaves, the morphism $\bar{\beta}(\Upsilon \times \Upsilon)$, which is the analogous morphism to $\Phi \times \Phi$ defined in subsection 3.3. We again denote this morphism by $\bar{\beta}$.

From the last Lemma and Lemma 4.9 we obtain:

Lemma 4.15. If

$$[(\bar{E}_j, \bar{i}_j, \bar{\tau}, g_{p+D}), (E_j, i_j, \tau, f_{p+D})] \in \bar{\beta}^{-1}(Z_n^{p+D})$$

defined over a field K , then:

1) There exists a $p+D$ -isogeny between the elliptic sheaves with $p+D$ -level structures $(\bar{E}_j, \bar{i}_j, \bar{\tau}, g_{p+D})$ and $(E_j, i_j, \tau, f_{p+D})$.

or

2) There exists an isogeny between the elliptic sheaf $(\bar{E}_j, \bar{i}_j, \bar{\tau})$ and the twisted elliptic sheaf $(E_j, i_j, \tau) \otimes_{\mathcal{O}_C} \mathcal{O}_C(-p-D)$.

or

3) $(F^\# \bar{E}_0)^\vee \otimes_{\mathcal{O}_C \otimes K} E_0((g+d)\infty - p - D) \in \Theta_{n^2}$.

Proof. Let consider us the diagram (not necessarily commutative)

$$\begin{array}{ccccc}
 \bar{E}_n & \xrightarrow{T(1)} & E_n(g+d-1) \\
 \bar{l} \uparrow & \searrow & \swarrow \uparrow l \\
 \bar{E}_1 & & (\mathcal{O}_{p+D})^n \otimes R & & E_1(g+d-1) \\
 \bar{\tau} \uparrow & \nearrow & \nwarrow \uparrow \tau \\
 F^\# \bar{E}_0 & \xrightarrow{F^\# T} & F^\# E_0(g+d-1)
 \end{array}$$

T is a morphism given by $[(\bar{E}_j, \bar{i}_j, \bar{\tau}, g_{p+D}), (E_j, i_j, \tau, f_{p+D})] \in \bar{\beta}^{-1}(Z_n^{p+D})$. The diagonal morphisms are the level structures, and the vertical morphisms $\bar{l}$ and l are induced by the " i_j " and " $\bar{i}_j$ ". From Definition 4.1, we have two possibilities: either $T: \bar{E}_0 \rightarrow E_0(g+d-1)$ takes values in $E_0(g+d-1)(-p-D)$ or the latter diagram is commutative after tensoring by $\mathcal{O}_{pD}$. In both cases, one obtains a morphism:

$$T(1) \cdot \bar{l} \cdot \bar{\tau} - l \cdot \tau \cdot F^\# T : F^\# \bar{E}_0 \rightarrow E_0((g+d)\infty - p - D).$$

When $T \cdot \bar{l} \cdot \tau \cdot t - \tau' \cdot l' \cdot t' \cdot F^\# T = 0$, we obtain cases 1) and 2) and when it is $\neq 0$ it tells us that $(F^\# \bar{E}_0)^\vee \otimes_{\mathcal{O}_C \otimes K} E_0((g+d)\infty - p - D) \in \Theta_{n^2}$. $\square$

Thanks to [An2] Lemma 3.3, for $n = 1$ it is possible to complete 3) in terms of Drinfeld modules. It would be very interesting to study this Lemma for rank- n -shtukas and rank- n -Drinfeld modules. For $C = \mathbb{P}_1$, because E_0 and $\bar{E}_0$ are semi-stable of 0 degree we have:

$$(F^\# \bar{E}_0)^\vee \otimes_{\mathcal{O}_C \otimes K} E_0((g+d)\infty - p - D) \notin \Theta_{n^2}.$$

One can study explicit examples, as in section 3, merely by changing monic polynomials " $q(t)$ " for polynomials $q(t)$ with $q(\alpha_p) = 1$, where $m_p = (t - \alpha_p) \cdot \mathbb{F}_q[t]$.

Remark 16. Given a rank- n -elliptic sheaf (E_j, i_j, τ) , we have that

$$h^0(E_{-n}) = h^1(E_{-n}) = 0.$$

Thus, E_{-n} is a semistable vector bundle. In this way, there exists a $d \gg 0$ such that for every pair of elliptic sheaves $[(\bar{E}_j, \bar{i}_j, \bar{\tau}), (E_j, i_j, \tau)]$ defined over a ring R is

$$H^1(C \otimes R, \bar{E}_0^\vee \otimes_{\mathcal{O}_C \otimes R} E_0((g+d-1))) = 0.$$

Therefore, $[(\bar{E}_j, \bar{i}_j, \bar{\tau}), (E_j, i_j, \tau)] \in \mathcal{U}_d$.

In the case of shtukas, this result is not true unless one considers shtukas with truncation [La1] and [La2]. In a following work, it would be interesting to study this issue in the setting of Lafforgue's shtuka compactifications [La1].

Acknowledgments I began to study this matter after a stay by Prof.G.W.Anderson in May 2002 at the University of Salamanca. The conversations, suggestions and the encouragement of Prof. G.W.Anderson have been invaluable to me for studying this issue in the rank- n case. I should like to express my gratitude to Ricardo Alonso for his friendship and company at all times and Jesus Muoz Diaz and Laurent Lafforgue for their friendship, desinterested help and Knidness towards me.

REFERENCES

- [Al] Alvarez, A. "Uniformizers for elliptic sheaves", International Journal of Mathematics, **11** n 7 (2000), 949-968
- [An1] Anderson, G. "t-motives", Duke Mathematical Journal. **53** (1986), pp.457-502
- [An2] Anderson, G. "Rank one elliptic modules A -modules and A -harmonic series", Duke Mathematical Journal. **73** (1994), pp.491-542
- [An3] Anderson, G. "A two dimensional analogue of Stickelberg's theorem " in: The Arithmetic of Function Fields, ed. D.Goss, D.R. Hayes, W. de Gruyter, Berlin, 1992, pp.51-77.
- [An4] Anderson, G. "Catalan transforms". Notes.
- [AnDP] Anderson, G. Dale Brownawell, W Papanikolas, M.A "Determination of the algebraic relations among special Γ -values in positive characteristic" alg-geom 0207168.
- [BlSt] Blum, A. Stuhler, U. "Drinfeld modules and elliptic sheaves", Vector bundles on curves-new directions (eds. S.Kumar et al.) LNM 1649 (1997)
- [DN] Drezet, J-M. Narasimhan, M.S. "Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques", Invent.Math. **97** (1989), pp.53-94
- [Dr1] Drinfeld, V.G. "Elliptic modules", English Transl.in Math.U.S.S.R-Sb. **23** n 4 (1976). pp.561-592
- [Dr2] Drinfeld, V.G. "Commutative subrings of certain non-commutative rings", English Transl. in funct.Anal.Appli. **21** (1987), pp.107-122
- [Dr3] Drinfeld, V.G. "Varieties of modules of F -sheaves", English Transl. in funct.Anal.Appli. **11** (1977), pp.9-12
- [C] Coleman, R. "On the Frobenius endomorphisms of the Fermat and Artin-Schreier curves" in: The Arithmetic of Function Fields, Proc. Amer. Math Soc, **102** (1988), pp.463-466.
- [La1] Lafforgue, L."Chtoucas de Drinfeld et correspondance de langlands" Invent.Math. n 1 **147** (2002), pp.1-241
- [La2] Lafforgue, L."Une compactification des champs classifiant les chtoucas de Drinfeld" Journal of the American Mathematical Society, (11) n 4, pp.1001-1036
- [L] Laumon, G. "Cohomology of Drinfeld Modular varieties", Cambridge University Press. **41** (1996)
- [LM] Laumon, G. Moret-Bailly, L. "Champs algébriques", Ergebnisse der Mathematik 39 **873** (1999)
- [LRSt] Laumon, G. Rapoport, M. Stuhler, U. "D-elliptic sheaves and the Langland's correspondence", Inventiones Mathematicae. **113** (1993), pp.217-338
- [Mu] Mumford, D. "An algebro-geometric construction of commuting operators and of solutions to the Toda lattice equations, Korteweg-de Vries equation and related non-linear equations ", Int.Symp. Algebraic Geometry (Kyoto 1977), Kinokuniya, Tokyo 1977, 115-153.
- [Se] Serre, J.P. "Groupes algébriques et corps de classes", Hermann (1959)
- [Ss] Seshadri, C.S. "Fibrés vectoriels sur les courbes algébriques", (rédigé par J-Drezet) Astérisque. **96** SMF(1982)
- [Si] Sinha, S. "Periods of t-motives and transcendence", Duke Mathematical Journal. **88** (1997), pp.465-535.

Álvarez Vázquez, Arturo
e-mail: aalvarez@gugu.usal.es

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE SALAMANCA, PLAZA DE LA MERCED 1-4.
 SALAMANCA (37008). SPAIN.